

# An Exception to the Second Law of Thermodynamics

**Yongle Yu**

State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics and Mathematics, Chinese Academy of Science, West No. 30 Xiao Hong Shan, Wuchang, Wuhan, 430071, China

E-mail: [yongle.yu@wipm.ac.cn](mailto:yongle.yu@wipm.ac.cn)

**Abstract.** We show that an exception to the second law of thermodynamics can be made by using superfluid  $^4\text{He}$ .

PACS numbers: 05.70.-a, 67.25.dg, 67.40.Kh

*Keywords:* superflow, entropy, many-body spectra

*Date:* July 20, 2106

The second law of thermodynamics (SLT), ruling out the decrease of the entropy in an isolated system, was first formulated by Clausius in 1850s and is regarded as a cornerstone of physics. Despite more than one and a half century of studies, it remains elusive to find a solid principle behind SLT. This could suggest that SLT is almost universal rather than exactly universal, and that there is a few exceptions to SLT. On the other hand, an exception to SLT could open up the possibility of rendering usable the thermal energy of materials in the environment, thus solving the energy issues faced by modern civilization. In this note, we show that an exception to SLT can be made by using superfluid  $^4\text{He}$ , and that it is possible to build a system for extracting thermal energy from the environment.

The phenomenological two-fluid model of superfluid  $^4\text{He}$  postulates that a superflow carries no heat or entropy. This postulation is not corroborated by any microscopic understandings of superfluidity. On the contrary, a recently developed theory [1], built in the natural quantum mechanics terms of many-body quantum states and jumps among the states, suggests that a superflow carries heat. This microscopic theory reveals that the many-body spectrum of a superfluid has a salient feature [1, 2, 3].


If one schematically plots the energy levels of a superfluid at the energy-momentum ( $E - P$ ) plane (see Figure 1), one then spots that the low-lying levels with various momenta form valley-like structures at the boundary region. At low temperature, only those levels within the bottoms of the valleys are relevant in physics. A superflow state corresponds to some occupied levels at a valley at a non-zero momentum. The system can't jump from the occupied levels at this valley to unoccupied levels at the left-side neighboring valley, due to the energy barrier which separates the valleys. The prohibition of inter-valley jumps ensures that a superflow keeps its momentum, thus causing the phenomenon of superfluidity [4].

Under this microscopic description, a  ${}^4\text{He}$  superflow shall always exchange quanta with its surroundings, as a result of the microscopic interactions between the  ${}^4\text{He}$  atoms in the superflow and the particles in the surroundings. This exchange process promotes the superflow's intra-valley jumps among the occupied low-lying levels at the corresponding valley. It leads to a thermal equilibrium occupation probability of the levels, namely, the occupation probability of a level at the valley is proportional to  $e^{-E/kT}$ , where  $E$  is the energy of the level,  $k$  is the Boltzmann constant and  $T$  is the temperature of the superflow. This exchange process also establishes a thermal equilibrium between the superflow and its surroundings, in the sense that the temperature of the superflow is the same as the temperature of its surroundings.

Clearly, this natural framework shows not only that a superflow carries heat, but also that superfluidity doesn't require a superflow with zero entropy. One could also note that, from a general physics viewpoint, a macroscopic flow with zero entropy, immune from thermal influence of its surroundings, is rather too peculiar to be understood.

The heat content of a superflow is in principle determined by its physical settings, such as its geometry, its temperature and its flow velocity. Let us consider a  ${}^4\text{He}$  superflow passing through a channel where second half of the channel is empty but the first half is packed with a medium such as fine powders. If one keeps the temperature of the superflow the same when it exits the medium, one could expect that the heat content of the superflow shall change due to the change of its physical settings. If the change of heat content is negative, for the conservation of energy, one should provide heating to the superflow to maintain its temperature. Without such a heating, the temperature of the superflow shall drop when it enters the second half of the channel.

Shortly after the discovery of superfluidity [5], the temperature drops of  ${}^4\text{He}$  superflows, when passing from media to rather free space, were reported [6]. Some observations involved a superflow in a capillary or a film superflow. For convenience, we enhance the meaning of media so that superflows in various media also cover a superflow in a capillary and a film superflow. Free space is referred to as a medium of vacuum. Then one would expect a change of the temperature of a superflow going from a medium to a different medium, for the reasons illustrated above. This is analogous to Peltier effect, in which the heat content of an electric current shall change when it passes from one wire to another wire made of a different kind of metal, causing a cooling or heating at the junction of two wires.



**Figure 1.** A schematic plot of many-body eigen levels of a superfluid in the  $E - P$  Plane. Dots (in gray and in black) represent levels. The horizontal and vertical coordinate of a dot corresponds to the momentum and the energy of the level, respectively.  $\Delta$  denotes the height of an energy barrier between two "valleys". Dots in black are the occupied levels of the system (at low temperature), corresponding to a superflow state with a momentum of  $P_c$ .

We shall consider a  ${}^4\text{He}$  superflow running inside a torus-shaped vessel, where half of the vessel is packed with one kind of medium and the other half is packed with a different medium [7]. We refer to this system as a heterogeneous superflow loop (HSL). If an isolated HSL is waited for a sufficient time, one can reach an interesting temperature configuration along the system. Apart from the interfaces between two media, one part of the system has a temperature  $T_h$  higher than the temperature of the opposite part of the system (denoted by  $T_l$ ). At this configuration, the heat content of superflow in one medium is (approximately) the same as the heat content of superflow in the other medium. To demonstrate an exception to SLT, let us make a good thermal contact between the high temperature part of the system and an infinitely large heat reservoir at a temperature  $T_h^r$ , and make a good thermal contact between the low

temperature part of HSL to another infinitely large heat reservoir at a temperature  $T_l^r$ . Let  $T_h > T_h^r > T_l^r > T_l$ . Then, the superflow at the high temperature medium is cooled, thus carrying less heat content while the superflow at the low temperature medium is warmed and carries more heat content. Thus the superflow in the high temperature medium carries less heat content than the superflow in the low temperature medium. As a result, the superflow will transfer heat from the reservoir at  $T_l^r$  to the reservoir which has a higher temperature ( $T_h^r > T_l^r$ ). This is a process in which the entropy of the whole system decreases rather than increases. The entropy-decreasing process can run for a cosmologically long time, for that the decay of the superflow is prevented by some energy barriers [8].

It is rather straightforward to sketch out a system for extracting thermal energy from the environment. It shall include a HSL, a cryogenic heat device (CHD), another heat device referred to as the major heat device (MHD). The operation of each heat device shall generate a net effect similar to that of a heat engine; it absorbs heat from a hot object, converts part of the heat into another form of energy and releases the rest to a cold object. The CHD operates between the high temperature part and the low temperature part of the HSL. It converts part of heat, absorbed from the high temperature part of the HSL, into another form of energy and releases this energy to the environment [9]. The CHD passes some heat to low temperature part of the HSL, the temperature of this part changes accordingly and the superflow transfers heat back to the high temperature part. The MHD operates between a room temperature material and the high temperature part of the HSL. It absorbs heat from the material and converts part of the heat into usable energy or works. It is clear that MHD is responsible for generating the bulk of useful energy or work while the CHD mainly plays the role of returning the heat received by the HSL back to the environment. To improve the operation conditions for the CHD, one could use a series of HSLs for generating a large temperate difference across the series. In the series, the low temperature part of the first HSL is in good thermal contact with the high temperature part of the second HSL, the low temperature part of the second HSL in contact with the high temperature part of the third HSL, and so on.

Vacuum systems shall be involved to prevent the direct heat transfer, mediated by air, from the environment to the HSL. Thermal radiation from the environment to the HSL can be technically reduced but can not be effectively eliminated. If the thermal radiation upsets the operation conditions for the HSL, one can add a layer to surround the HSL. The MHD then operates between the room temperature material and the layer, and an additional cryogenic heat device works between the layer and the HSL. This layer has an intermediate temperature so that thermal radiation received by HSL can be effectively weakened. If needed or preferred, more layers can be implemented.

It is interesting to ask whether one can build a similar system based on superconductors instead of superfluid  $^4\text{He}$ . It seems that the electric supercurrents in many superconductors have too low heat contents to generate a realistic temperature difference along a similar heterogeneous supercurrent loop. However, in some heavy-

fermion superconductors, such as  $CeCoIn_5$  [10], the supercurrents might have significant workable heat contents .

In conclusion, we show that there is an exception to the second law of thermodynamics, and that this exception could lead to a solution to the energy issues.

Financial supports from Chinese NSF (Grant No. 11474313), from CAS (Grant No. XDB21030300), from NKRD of China ( Grant No. 2016YFA0301503), are acknowledged.

## References

- [1] Y. Yu, Ann. Phys. **323**, 2367 (2008), and references therein; Y. Yu, Mod. Phys. Lett. **B 29**, 1550068-1 (2015).
- [2] F. Bloch, Phys. Rev. **A 7**, 2187 (1973).
- [3] A. G. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
- [4] More generally, a superflow state could corresponds to a number of valleys with each of these valleys being partially occupied ( The occupation probability of one valley refers to the sum of the occupation probabilities of the levels at the valley). Since the occupation probability of each valley doesn't change (with time) due to the prohibition of inter-valley jumps, the superflow state will always keep its momentum.
- [5] P. Kaptiza, Nature **141**, 74 (1938); J. F. Allen and A. D. Misener, Nature **141**, 75 (1938) .
- [6] see, e. g., J. G. Daunt and K. Mendelsohn, Nature, **143**, 719 (1939).
- [7] Here a medium could also refer to a vacuum, and to a specific geometric setting of a superflow.
- [8] For experimental observations of persistent currents of liquid  $^4\text{He}$ , see, e. g., J. D. Reppy and D. Depatie, Phys. Rev. Lett. **12**, 187 (1964).
- [9] For example, one can use a number of thermocouples between the high temperature part and the low temperature part of the HSL. these thermocouples, connected in series and in parallel, could generate a workable voltage and current. If preferred, the voltage and current can be passed without dissipation to a separate cryogenic device placed away, by using superconducting wires. This cryogenic device could work at a temperature of several tens of kelvins for the sake of its performance (The heat leak through the superconducting wires will be negligible if the wires were sufficient long). This device could be a number of light-emitting diodes to release energy in the form of photons, into the environment. Or it could convert the (direct) current to alternating current, which in turn drives a superconducting solenoid to generate an oscillating magnetic field. The magnetic energy then can be picked up by a room-temperature solenoid outside, which is physically not in touch with the cryogenic parts. One might also develop cryogenic thermomagnetic generators for this use.
- [10] C. Petrovic, P. G. Pagliuso, M. F. Hundley, R. Movshovich, J. L. Sarrao, J. D. Thompson, Z. Fisk and P. Monthoux, J. Phys. Condens. Matter **13**, L337 (2001).