# Geochemical signatures and human health risk evaluation of rare earth elements in soils and plants of the northeastern Qinghai-Tibet Plateau, China

LI Leiming<sup>1</sup>, WU Jun<sup>2\*</sup>, LU Jian<sup>3,4</sup>, ZHANG Xiying<sup>1,5</sup>, XU Juan<sup>6</sup>

<sup>1</sup> Key Laboratory of Comprehensive and Highly Efficient Utilization of Salt Lake Resources, Qinghai Institute of Salt Lakes, Chinese Academy of Sciences, Xining 810008, China;

<sup>2</sup> Yantai Research Institute, Harbin Engineering University, Yantai 264006, China;

<sup>3</sup> CAS Key Laboratory of Coastal Environmental Processes and Ecological Remediation, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China;

<sup>4</sup> Shandong Key Laboratory of Coastal Environmental Processes, Yantai 264003, China;

<sup>5</sup> Qinghai Provincial Key Laboratory of Geology and Environment of Salt Lakes, Xining 810008, China;

<sup>6</sup> State Key Laboratory of Marine Geology, Tongji University, Shanghai 200092, China

Abstract: Information on rare earth elements (REEs) in soils and plants of the Qinghai-Tibet Plateau is very limited. Therefore, in this study, we performed field sampling to explore the geochemical signatures and human health risk of REEs in soils and plants of the northeastern Qinghai-Tibet Plateau, China. A total of 127 soil samples and 127 plant samples were collected from the northeastern Qinghai-Tibet Plateau to acquire the geochemical signatures and related human health risks of REEs. The mean total concentrations of REEs in soils and plants of the study area reached 178.55 and 10.06 mg/kg, respectively. The light REEs in soils and plants accounted for 76% and 77% of the total REEs, respectively. REEs showed significantly homogenous distribution in soils but inhomogeneous distribution in plants of the study area. Characteristic parameters indicated that light REEs were enriched and fractionated significantly, while heavy REEs were moderately fractionated in soils and plants. REEs in soils and plants showed significantly negative Europium anomaly. Cerium showed slightly positive anomaly in plants and slight anomaly in soils. The normalized distribution patterns of REEs were generally similar in the analyzed soils and the corresponding plants of the study area. The average bio-concentration factor of REEs ranged from 0.0478 (Scandium) to 0.0604 (Europium), confirming a small accumulation of REEs by plants. Health risks caused by REEs in soils and plants were negligible, while risks for adults were lower than those for children. This study provides important information on REEs in soils and plants of the northeastern Qinghai-Tibet Plateau.

Keywords: rare earth elements; geochemical signatures; human health risk; carcinogenic risk; bio-concentration factor; Qinghai-Tibet Plateau

**Citation:** LI Leiming, WU Jun, LU Jian, ZHANG Xiying, XU Juan. 2022. Geochemical signatures and human health risk evaluation of rare earth elements in soils and plants of the northeastern Qinghai-Tibet Plateau, China. Journal of Arid Land, 14(11): 1258–1273. https://doi.org/10.1007/s40333-022-0107-8

# **1** Introduction

Rare earth elements (REEs) encompass the yttrium (Y), the elements of the lanthanide series (atomic number from 57 to 71), and scandium (Sc), which exhibit similar environmental

<sup>\*</sup>Corresponding author: WU Jun (E-mail: wujunlisa@163.com)

Received 2022-07-06; revised 2022-10-09; accepted 2022-10-14

<sup>©</sup> Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Science Press and Springer-Verlag GmbH Germany, part of Springer Nature 2022

behaviors (Ferreira et al., 2021). Promethium (Pm) does not exist in the Earth's crust since the radioactive decay of Pm is fast (Migaszewski and Gałuszka, 2015). REEs generally exist together in natural environment due to similar physical and chemical properties (Groenenberg et al., 2010; Khan et al., 2017). Many studies reported that pedogenic processes, soil characteristics, and parent material affected REEs background contents in soils (Yao et al., 2010; Wu et al., 2019). REEs are not rare from a geochemical perspective because the abundances of REEs in soils are similar to those of zinc (Zn), lead (Pb), and copper (Cu) and higher than those of mercury (Hg), Stannum (Sn), cobalt (Co), and silver (Ag) (Wang et al., 1998).

Heavy REEs ( $\Sigma$ HREEs; the sum Y, Sc, and from gadolinium (Gd) to lutetium (Lu)) and light REEs ( $\Sigma$ LREEs; the sum from lanthanum (La) to europium (Eu)) compose REEs based on the mass and atomic numbers of REEs. The concentrations of  $\Sigma$ HREEs are generally lower than the concentrations of  $\Sigma$ LREEs (Henderson, 1984; Durn et al., 2021). The  $\Sigma$ LREEs are more mobile and available for plant absorption than  $\Sigma$ HREEs (Brioschi et al., 2013). The concentrations of  $\Sigma$ LREEs in leachate are higher than those of  $\Sigma$ HREEs so that the  $\Sigma$ LREEs are more available for root adsorption than  $\Sigma$ HREEs (Aide and Aide, 2012). The bioavailability of REEs in soil is a considerable factor for the accumulation of REEs by plants (Li et al., 2013; Liu et al., 2018). The concept of bioavailability also was defined the amount of human adsorption in recent years (Li et al., 2018; Ferreira et al., 2022; Wang et al., 2022).

The global annual demands of REEs have increased significantly from  $3.00 \times 10^4$  to  $1.35 \times 10^5$  t during 1980-2015 (Barta, 2007; Huang et al., 2016). China is the largest producer of REEs in the world, accounting for approximately 97% of the global supply. The largest REEs ore mine is located in the northern China (Tang et al., 2020). REEs have been widely used in medical technology, industry, and agriculture (Hu et al., 2006; Pagano et al., 2015; Wang and Liang, 2016; Ou et al., 2022). Various field activities have released lots of REEs into the environment (Zhang et al., 2014; Pagano, 2016). Distribution, environmental behaviors, and removal of different contaminants, such as trace elements (Zheng et al. 2010a, b; Liang et al., 2013; Rodríguez-Barranco et al., 2014; Lu et al., 2021; Pirarath et al., 2021; Cui et al., 2022; He et al., 2022), antibiotics and corresponding resistance genes, perfluorooctane sulfonate, polycyclic aromatic hydrocarbons, and microplastics, have attracted attention widely (Liu et al., 2019; Fu et al., 2021; Qiao et al., 2021; Lu et al., 2022a, b). However, studies regarding the negative effect of REEs on natural ecosystems and humans are limited. REEs accumulate in various environmental media (including soil, atmosphere, and water body) after entering into the environment, and finally influence ecosystem via human activities or human health via REEs accumulation in the human bodies (Mihajlovic et al., 2015; Pagano, 2016; Wang et al., 2017; Wang et al., 2019; Godwyn-Paulson et al., 2022). The behavior of REEs in natural environment is strongly affected by soil redox conditions, organic matter content, pH, weathering processes, climate conditions, and parent rock (Zhang et al., 2014; Pagano, 2016; Allajbeu et al., 2016; Huang et al., 2019). There are three ways including ingestion, dermal adsorption, and inhalation for entering the human body (Meryem et al., 2016; Guo et al., 2019). Previous studies have proven that REEs accumulated in the human body can cause numerous diseases (Zhang et al., 2000; Zhao et al., 2017; Ferreira et al., 2022).

Soil is an important medium for receiving pollutants and nutrients to play an important role in ecosystems. In recent decades, the geochemical characteristics and health hazards of REEs have attracted much attention (Kan et al., 2017; Huang et al., 2019; Guo et al., 2019; Wu et al., 2019; Malhotra et al., 2020). However, information about distribution patterns, accumulation in soils and plants, and health risk evaluation of REEs is quite scarce in the study area. The main objectives of this study were to (1) investigate the concentration and distribution patterns of REEs in soils and plants of the northeastern Qinghai-Tibet Plateau; (2) calculate bioavailability and mobility characteristics of REEs; and (3) evaluate the potential human health risk of REEs.

# 2 Materials and methods

# 2.1 Study area and soil and plant samples

The study area is situated in the northeastern Qinghai-Tibet Plateau of China, which spans  $4.13 \times 10^5$  km<sup>2</sup> (90°37′-101°05′E, 35°01′-38°43′N). The study area has an average annual precipitation of 50–450 mm and an annual average temperature of -5.7-8.5°C. A total of 127 soil samples (soil depth: 0–20 cm) and 127 plant samples were collected from June 14 to June 29 of 2017 by following standard sampling method (State Environmental Protection Administration, 2004) (Fig. 1). All samples were placed in plastic bags, transported back to the laboratory, and stored at  $-80^{\circ}$ C. Soil samples were prepared by following the study conducted by Li et al. (2018).



Fig. 1 Study area and soil and plant sampling sites

# 2.2 Analysis of rare earth elements (REEs)

The soil samples were air-dried, crushed, and then sieved using 0.074-mm sieves. The soil samples were calcined at 700°C for 2 h to eliminate the organic matter. According to related standard (State Environmental Protection Administration, 2004), 100 mg of each soil power was added along with 3 mL HNO<sub>3</sub> with high purity, 2 mL HCl with high purity, and 2 mL HF with high purity in a 100-ml Teflon digestion tube (Wu et al., 2019). These tubes were placed in a microwave dissolving system for 1 h and cooled. The cooled solution was dried at 180°C with 2 mL of high quality HClO<sub>4</sub> and 15 mL of high purity HNO<sub>3</sub>.

The plant (*Potentilla anserina* L.) samples were pre-treated by following the previously-reported procedures (Shen et al., 2018). All parts of plant samples were washed thoroughly to remove soil particles, washed five times with ultra-pure water, blotted extra water with tissue paper, and dried. Then the samples were grinded to fine powder for digestion. Approximately 100 mg of dry plant powder was digested with 3 mL HCl (37%), 5 mL HNO<sub>3</sub> (65%), and 1 mL HF (65%) (Ayrault et al., 2001; Li et al., 2018). The other digestion procedures for plant samples were same as soils. Using testing instrument (Agilent7900, Agilent Inc, Santa Clara, USA), we measured the concentration levels of La, cerium (Ce), praseodymium (Pr), neodymium (Nd), samarium (Sm), Eu, Gd, terbium (Tb), dysprosium (Dy), holmium (Ho), erbium (Er), thulium (Tm) ytterbium (Yb), Lu, Y and Sc in samples.

# 2.3 Distribution characteristics of REEs

North American shale composite (NASC) was used to normalize REEs concentrations to eliminate the "Oddo-Harkin effect" of REEs (Taylor and Mclennan, 1985). Normalization values

were calculated by the ratio of the REEs concentrations to the corresponding element concentrations of NASC in this study. The parity effects of REEs were eliminated after the normalized REEs by NASC (Chen and Yang, 2010; Wu et al., 2019).

Characteristic parameters which indicated the distribution of REEs in soils and plants included fractionation between  $\Sigma$ LREEs and  $\Sigma$ HREEs ((La/Yb)<sub>N</sub>), fractionation between  $\Sigma$ LREEs ((La/Sm)<sub>N</sub>), fractionation between  $\Sigma$ HREEs ((Gd/Yb)<sub>N</sub>), Ce anomaly ( $\delta_{Ce}$ ), and Eu anomaly ( $\delta_{Eu}$ ); we calculated these parameters based on normalized values of NASC (Taylor and Mclennan,1985; Mclennan, 1989; Chen and Yang, 2010; Wu et al., 2019; Yu et al., 2019), the calculation process is as follows:

$$(\text{La/Sm})_{\text{N}} = \frac{(\text{La}_{\text{sample}}/\text{La}_{\text{NASC}})}{(\text{Sm}_{\text{sample}}/\text{Sm}_{\text{NASC}})},$$
(1)

$$(Gd/Yb)_{N} = \frac{(Gd_{sample}/Gd_{NASC})}{(Yb_{sample}/Yb_{NASC})},$$
(2)

$$(La/Yb)_{N} = \frac{(La_{sample}/La_{NASC})}{(Yb_{sample}/Yb_{NASC})},$$
(3)

$$\delta_{Ce} = \frac{(Ce_{sample}/Ce_{NASC})}{\sqrt{(La_{sample}/La_{NASC}) \times (Pr_{sample}/Pr_{NASC})}},$$
(4)

$$\delta_{\rm Eu} = \frac{(La_{\rm sample}/La_{\rm NASC})}{\sqrt{(Sm_{\rm sample}/Sm_{\rm NASC})/(Gd_{\rm sample}/Gd_{\rm NASC})}},$$
(5)

where  $La_{sample}$  (mg/kg),  $Sm_{sample}$  (mg/kg),  $Gd_{sample}$  (mg/kg),  $Yb_{sample}$  (mg/kg),  $Ce_{sample}$  (mg/kg), and  $Pr_{sample}$  (mg/kg) are the contents of La, Sm, Gd, Yb, Ce, and Pr in soil or plant samples, respectively; and  $La_{NASC}$  (mg/kg),  $Sm_{NASC}$  (mg/kg),  $Gd_{NASC}$  (mg/kg),  $Yb_{NASC}$  (mg/kg),  $Ce_{NASC}$  (mg/kg), and  $Pr_{NASC}$  (mg/kg) are the contents of La, Sm, Gd, Yb, Ce, and Pr in reference material, respectively. The content of REEs in reference materials can be found in Taylor and Mclennan (1985).

#### 2.4 Bio-concentration factor (BCF)

Bio-concentration factor (BCF) is usually adopted to calculate the potential transfer of various contaminants from soils to plants (Liu et al., 2015; Ingrid et al., 2022; Nicolas et al., 2022). This approach was also employed to examine REEs. The BCF is calculated as follows:

$$BCF_i = \frac{X_{\text{plant}}}{X_{\text{soil}}},\tag{6}$$

where BCF<sub>*i*</sub> is the BCF value of the  $i^{\text{th}}$  target REEs; and  $X_{\text{plant}}$  (mg/kg) and  $X_{\text{soil}}$  (mg/kg) are the concentrations of the target REEs in soil and plant samples, respectively.

#### 2.5 Human health risk evaluation

Three exposure pathways including ingestion, inhalation, and dermal adsorption of REEs in soils and plants might exert threat to human health (Zheng et al., 2020). This study used formulas from the Environmental Protection Agency of USA and other articles (USEPA, 1996; Chen et al., 2022; Li et al. 2022). The hazard quotients posed through three ways can be calculated using the following equations:

$$HQ_{ing} = C \times \frac{IngR \times EF \times ED}{BW \times AT \times CFD} \times 10^{-6},$$
(7)

$$HQ_{inh} = C \times \frac{InnR \times EF \times ED}{PEF \times BW \times AT \times CFD},$$
(8)

#### JOURNAL OF ARID LAND

$$HQ_{der} = C \times \frac{SAF \times SAE \times DAF \times EF \times ED}{BW \times AT \times CFD} \times 10^{-6}, \qquad (9)$$

$$HI = HQ_{ing} + HQ_{inh} + HQ_{der}, \qquad (10)$$

where HQ<sub>ing</sub>, HQ<sub>inh</sub>, and HQ<sub>der</sub> are the hazard quotient of ingestion, inhalation, and dermal adsorption, respectively; *C* is element concentration (mg/kg); BW is body weight (kg); EF is exposure frequency (d/a); ED is exposure duration (a); AT is averaging time (d); CFD is corresponding reference dose (mg/(kg-d)); IngR (mg/d) and InhR (m<sup>3</sup>/d) are ingestion rate and inhalation rate, respectively; PEF is particle emission factor (m<sup>3</sup>/mg); SAF is skin adherence factor (mg/(cm<sup>2</sup>·d)); SAE is skin area exposed (cm<sup>2</sup>); DAF is dermal absorption factor; and *HI* is the sum of HQ<sub>ing</sub>, HQ<sub>inh</sub>, and HQ<sub>der</sub>.

Reference values used in the equations are from previous studies (USEPA, 2002; Duan, 2012; Faiz et al., 2012; Chen et al., 2022; Li et al., 2022). We used a unique reference dose value (0.02 mg/(kg·d)) for all REEs in this study (Li et al., 2013; Sun et al., 2017). The values of hazard quotient above 1.0 are considered non-carcinogenic risk to human health and the values of hazard quotient below 1.0 are considered safe for the human body (Wu et al., 2018; Li et al., 2022). Carcinogenic risk was calculated by lifetime mean daily dose (LMDD, mg/(kg·d)) (Sun et al., 2017; USEPA, 2004) and used the following formulas:

$$LMDD = \frac{C \times EF}{AT} \times \left(\frac{CR_{child} \times ED_{child}}{BW_{child}} + \frac{CR_{adult} \times ED_{adult}}{BW_{adult}}\right),$$
(11)

$$Carcinogenic risk = LMDD \times SF,$$
(12)

where  $CR_{child}$  and  $CR_{adult}$  are contact rate for child and adult, respectively;  $ED_{child}$  and  $ED_{adult}$  are exposure duration for children and adults (a), respectively;  $BW_{child}$  and  $BW_{adult}$  are body weight of children and adults (kg), respectively; and SF is slope factor, which is  $3.2 \times 10^{-12}$  in the study (Sun et al., 2017). The unacceptable risk for regulatory purposes is higher than  $1.0 \times 10^{-4}$  (Faiz et al., 2012; Chen et al., 2022).

#### 2.6 Statistical analysis

The statistical data of REEs were obtained by Excel while the correlation of REEs with soils or plants was studied by Pearson correlation analysis by using SPSS 22.0. Digital terrain model was established by applying inverse distance weight interpolation (Li et al., 2013; Xu et al., 2016; Wang et al., 2019; Wu et al., 2021). ArcGIS 10.2 processed corresponding distribution for data on human health risk values.

#### **3** Results and discussion

## 3.1 REEs of soils

Detailed statistical results on the concentrations of total REEs and individual REEs in 127 soil samples were showed in Table 1. The concentrations of REEs normalized NASC, background values in Qinghai Province of China, and background values in China were presented (Taylor and Mclennan, 1985; CNEMC, 1989; MEPC, 1990). The concentrations of REEs in soils ranged from 55.29 to 306.80 mg/kg with an average of 178.55 mg/kg (Table 1). The concentration of REEs in soils from most of sampling sites were generally similar in the study area. The REEs exhibited significantly good homogeneity of distribution in soils. The total concentrations of REEs in 127 soil samples exceeded 100.00 mg/kg. Two soil samples with the highest concentrations of REEs were located in an industrial area (more than 300.00 mg/kg). The results indicated that anthropogenic activities, particularly mining or industrial activities, may contribute slightly to the accumulation of REEs in some sites of study area.

The average concentrations of REEs of NASC, background values in Qinghai Province of China, and background values in China were lower than the concentrations in soils of the study area (Taylor and Mclennan, 1985; CNEMC, 1990; MEPC, 1990). Likewise, the average concentrations

of  $\Sigma$ LREEs in soils (135.26 mg/kg) were lower than the background values in China (147.92 mg/kg) and higher than that of NASC (73.50 mg/kg) and background values in Qinghai Province of China (126.37 mg/kg). Mean concentrations of  $\Sigma$ HREEs (43.29 mg/kg) were higher than the  $\Sigma$ HREEs of NASC (33.40 mg/kg), background values in Qinghai Province of China (35.46 mg/kg), and background values in China (38.84 mg/kg) (Table1).  $\Sigma$ LREEs in soils accounted for 76% of the total REEs, which was consistent with Tyler's results (Tyler, 2004). The results indicated that concentrations of  $\Sigma$ LREEs were higher than those of  $\Sigma$ HREEs, similar to the results of NASC. The correlation analysis illustrated that there was an obvious positive correlation among REEs of soils (Fig. S1), while the relationship among Eu, Er, Sc, and other REEs was poor. These results indicated that most of REEs might have similar sources in the soils.

 Table 1
 Statistic parameters of rare earth elements (REEs) in soils and plants of the northeastern Qinghai-Tibet

 Plateau, China
 Plateau

		Soils (1	mg/kg)		Plants (	mg/kg)			Background	Background values in China (soils) <sup>c</sup> (mg/kg)	
Rare earth - element and parameter	Min Mean		Max	SD	Min	Mean	Max	SD	NASC <sup>a</sup> (mg/kg)		values in Qinghai Province of China (soils) <sup>b</sup> (mg/kg)
La	8.36	31.04	54.26	6.6	0.10	1.76	8.66	1.7	16.00	32.80	39.70
Ce	17.09	63.66	112.43	13.2	0.44	3.78	16.70	3.4	33.00	58.30	68.40
Pr	1.82	7.32	12.50	1.6	0.02	0.39	1.93	0.4	3.90	5.87	7.17
Nd	6.95	26.95	46.38	5.8	0.08	1.43	7.19	1.4	16.0	23.70	26.40
Sm	1.33	5.18	8.94	1.1	0.02	0.28	1.45	0.3	3.50	4.77	5.22
Eu	0.33	1.11	3.26	0.3	0.00	0.07	0.32	0.1	1.10	0.93	1.03
Gd	1.31	5.10	8.81	1.0	0.02	0.28	1.41	0.3	3.30	4.15	4.60
Tb	0.18	0.68	1.24	0.1	0.00	0.04	0.19	0.0	0.60	0.60	0.63
Dy	0.99	3.56	6.68	0.7	0.01	0.20	1.02	0.2	3.70	3.80	4.13
Ho	0.20	0.69	1.29	0.1	0.00	0.04	0.20	0.0	0.80	0.77	0.87
Er	0.64	2.19	6.17	0.7	0.01	0.11	0.58	0.1	2.20	2.32	2.54
Tm	0.08	0.28	0.51	0.1	0.00	0.02	0.08	0.0	0.30	0.33	0.37
Yb	0.53	1.82	3.32	0.4	0.01	0.11	0.52	0.1	2.20	2.07	2.44
Lu	0.08	0.27	0.47	0.1	0.00	0.02	0.08	0.0	0.30	0.32	0.36
Y	5.13	18.20	34.64	3.9	0.07	1.07	5.43	1.0	20.00	21.10	22.90
Sc	4.86	10.51	21.93	3.2	0.03	0.47	2.48	0.4	30.00	10.42	/
REEs	55.29	178.55	306.80	35.8	1.03	10.06	48.25	9.3	136.90	172.25	186.76
ΣLREEs	35.88	135.26	234.42	28.2	0.84	7.71	36.27	7.2	73.50	126.37	147.92
ΣHREEs	18.34	43.29	72.39	8.8	0.18	2.35	11.98	2.2	33.40	35.50	38.84
ΣLREEs/Σ HREEs	2.75	4.14	6.09	0.4	2.95	4.21	6.74	0.6	2.20	3.56	3.81
(La/Yb) <sub>N</sub>	1.71	2.36	3.61	0.3	1.52	2.22	3.01	0.3	/	2.18	2.24
(La/Sm) <sub>N</sub>	1.16	1.31	1.54	0.1	1.08	1.36	1.67	0.1	/	1.50	1.66
$(Gd/Yb)_N$	1.50	1.87	2.51	0.2	1.29	1.78	3.06	0.2	/	1.34	1.26
$\delta_{Ce}$	0.95	1.01	1.31	0.1	0.93	1.23	3.06	0.4	/	1.01	0.97
$\delta_{Eu}$	0.49	0.67	2.02	0.1	0.51	0.81	3.14	0.3	/	0.65	0.65

Note: La, lanthanum; Ce, cerium; Pr, praseodymium; Nd, neodymium; Sm, samarium; Eu, europium; Gd, gadolinium; Tb, terbium; Dy, dysprosium; Ho, holmium; Er, erbium; Tm, thulium; Yb, ytterbium; Lu, lutetium; Y, yttrium; Sc, scandium; REEs, rare earth elements;  $\Sigma LREEs$ , light REEs (the sum from La to Eu);  $\Sigma HREEs$ , Heavy REEs (the sum Y, Sc, and from Gd to Lu);  $(La/Yb)_N$ , fractionation between  $\Sigma LREEs$  and  $\Sigma HREEs$ ;  $(La/Sm)_N$ , fractionation between  $\Sigma LREEs$ ;  $(Gd/Yb)_N$ , fractionation between  $\Sigma HREEs$ ;  $\delta_{Ce}$ , Ce anomaly;  $\delta_{Eu}$ , Eu anomaly; Min, minimum; Max, maximum; SD, standard deviation; NASC, North American shale composite. <sup>a</sup> refers to Taylor and Mclennan (1985), <sup>b</sup> refers to MEPC (1990), <sup>c</sup> refers to CNEMC (1989), and / refers to no data.

#### 3.2 REEs of plants and BCF

Table 1 also showed the concentrations of total and individual REEs in plants. The order of concentrations of REEs in plants was basically the same as that in soils. The distribution of REEs in plants was extremely uneven in the study area. The average concentrations of REEs in plants were variable in different sites. The average concentration of total REEs was 10.06 mg/kg, and while the concentrations of total REEs ranged from 1.03 to 48.25 mg/kg in plants. The average concentration of  $\Sigma$ LREEs was 7.71 mg/kg in plants, and the concentration range of  $\Sigma$ LREEs was 0.84 to 36.27 mg/kg. The concentrations of  $\Sigma$ HREEs ranged from 0.18 to 11.98 mg/kg, with an average value of 2.35 mg/kg.  $\Sigma$ LREEs covered 77% of total REEs in plants of the study area. La, Ce, Nd, and Y in plants accounted for more than 80% of total REEs. The characteristics of REEs in soils of the study area were well correlated with the high fraction of the above REEs and the distribution order of REEs concentrations. The correlation analysis showed that there was a significantly positive correlation among the REEs in plants to prove that all REEs in plants came from the same sources (Fig. S1b).

BCF can often reflect a plant's capacity to accumulate certain components. The statistical information on BCF values of REEs at various sites was shown in Figure 2. At the sampling sites, BCF values ranged from 0.004 to 0.289 for all target REEs. The average BCF values for REEs followed the order of Eu>Lu>Y>Ce>Yb>Ho>Tm>Dy>La>Tb>Er>Gd>Sm>Nd>Pr>Sc. The average BCF value of Eu in the study area was high compared to other REEs, indicating a relatively high mobility of transplants in soils compared to other REEs. In the same study location, the mean BCF values of REEs were significantly lower than those of heavy metals (Li et al., 2018). However, the average BCF of REEs was lower than that of heavy metals, indicating that REEs were more difficult to transfer from soils to plants (Hu et al., 2013; Jeelani et al., 2017; Li et al., 2018). BCF values of REEs in all sites were less than 1.000, indicating relatively low bioaccumulation. The results were similar to the recent studies (Ingrid et al., 2022; Nicolas et al., 2022).



**Fig. 2** Bio-concentration factor (BCF) values of rare earth elements (REEs) in the study area. La, lanthanum; Ce, cerium; Pr, praseodymium; Nd, neodymium; Sm, samarium; Eu, europium; Gd, gadolinium; Tb, terbium; Dy, dysprosium; Y, yttrium; Ho, holmium; Er, erbium; Tm, thulium; Yb, ytterbium; Lu, lutetium; Sc, scandium. The boxes represent the range from the lower quantile (Q25) to the upper quantile (Q75). The dots and horizontal lines inside the boxes represent the means and medians, respectively. The dots outside the boxes represent outliers. The upper and lower whiskers show the range within 1.5IQR (interquartile range).

## 3.3 REEs Characteristic parameter in soils and plants

Table 1 and Figure 3 showed the characteristic parameters of REEs in soils and plants, including  $\delta_{Eu}$ ,  $\delta_{Ce}$ ,  $\Sigma LREEs/\Sigma HREEs$ ,  $(La/Sm)_N$ ,  $(Gd/Yb)_N$ , and  $(La/Yb)_N$ . The average value of  $\Sigma LREEs/\Sigma HREEs$  in soils was 4.14, and the range of  $\Sigma LREEs/\Sigma HREEs$  in soils was 2.75–6.09. The results were similar to previous report (Wu et al., 2018). The range of  $\Sigma LREEs/\Sigma HERRs$  in plants was 2.95–6.74 with an average value of 4.21, indicating that the values of  $\Sigma LREEs$  were more abundant in plants than soils. The characteristic parameter of  $(La/Sm)_N$  reflected the fractionation degree of  $\Sigma LREEs$ . The average value of  $(La/Sm)_N$  in soils was 1.31, while the range of  $(La/Sm)_N$  values was 1.16–1.54 in soils. The average value of  $(La/Sm)_N$  in plants was 1.36, while the range of  $(La/Sm)_N$  values was 1.08–1.67. These results indicated that  $\Sigma LREEs$  in soils and plants were significantly fractionated in the study area.



**Fig. 3** Concentrations of light REEs ( $\Sigma$ LREEs) and heavy REEs ( $\Sigma$ HREEs) as well as the ratio of  $\Sigma$ LREEs to  $\Sigma$ HREEs ( $\Sigma$ LREEs/ $\Sigma$ HREEs) in soil samples (a) and plant samples (b). P-1–P-127 represent the serial number of sampling sites.

The characteristic parameter of  $(Gd/Yb)_N$  illustrated the fractionation degree of  $\Sigma$ HREEs. The range of  $(Gd/Yb)_N$  values in soils was 1.50–2.51 with an average value of 1.87, while  $(Gd/Yb)_N$  values in plants ranged from 1.29 to 3.06 with an average value of 1.78 in the study area, illustrating that  $\Sigma$ HREEs in soils and plants had moderate fractionation. The characteristic parameter of  $(La/Yb)_N$  showed the fractionation degree of  $\Sigma$ LREEs and  $\Sigma$ HREEs. The range of  $(La/Yb)_N$  values in soils ranged from 1.71 to 3.61 with an average value of 2.36, while the range

of  $(La/Yb)_N$  values in plants was 1.52–3.01 with an average value of 2.22, illustrating that  $\Sigma LREEs$ -enriched and the curve of the figure was right oblique (Fig. 4). These characteristic parameters in this study were similar to the background values in Qinghai Province of China and the background values in China.



**Fig. 4** (a), normalization of average REEs values and background composition in soils; (b), normalization of average REEs values in plants. NASC, North American shale composite; BC, background composition. Soil/NASC represents normalization of average REEs values in soils, BC/NASC denotes normalization of background composition in soils, and Plant/NASC represents normalization of average REEs values in plants.

 $\delta_{Eu}$  and  $\delta_{Ce}$  generally indicate anomalies of Eu and Ce (Yao et al., 2010; Gill et al. 2018). The elements are positive anomalies when  $\delta_{Eu}$  or  $\delta_{Ce}$  values are greater than 1.05, conversely, they are negative anomalies when  $\delta_{Eu}$  or  $\delta_{Ce}$  values are lower than 0.95 (Zhao et al. 2017; Gill et al. 2018; Wu et al., 2019). The range of  $\delta_{Eu}$  values in soils was 0.49–2.02 with an average value of 0.67 in the study area. The average  $\delta_{Eu}$  value in plants was 0.81, ranging from 0.51 to 3.14. These results indicated Eu in soils and plants of the study area was significantly negative anomalies. The range of  $\delta_{Ce}$  values in soils was 0.95–1.31 with an average value of 1.01. The range of  $\delta_{Ce}$  values in plants was 0.93–3.06 with an average value of 1.23. These results illustrated that Ce did not show significantly anomaly in soils while it showed slightly positive anomaly in plants. The  $\delta_{Eu}$  and  $\delta_{Ce}$  of REEs in soils and plants of the study area were very similar to the background values in Qinghai Province of China and the background values in China.

Because REEs with even atomic numbers are always more numerous than those with odd atomic numbers, it is challenging to compare the abundance of REEs directly. The concentrations of REEs were normalized using NASC to remove "Oddo-Harkin effect" in the study area (Fig. 4). The normalization model can be adopted to compare the REEs abundance in various environmental media and to determine whether there is a shortfall or enrichment of REEs (Wu et al., 2019; Ferreira et al., 2021; Lin et al., 2021). The NASC-normalized distribution patterns of REEs were usually similar in soils and plants. The average REEs patterns of the soils and plants matched the average values patterns of the background values in Qinghai Province of China (Fig. 4a and b). The distribution patterns of REEs in soils and plants of the study area were similar to the REEs patterns of the sediments and soils (Wu et al., 2019; Costa et al., 2021). Similar distribution patterns of REEs demonstrated that they primarily came from natural sources.

#### 3.4 Health risk of soils and plants

Non-carcinogenic effects of REEs in soils and plants of the study area on human beings were calculated (Fig. 5; Tables S1 and S2). The total non-carcinogenic risks of REEs in soils for adults ranged from 0.0081 to 0.0450, while those in plants for adults ranged from 0.0002 to 0.0071. The range of total non-carcinogenic risks from REEs in soils for children was 0.0187 to 0.1038, while

the total non-carcinogenic risks of REEs in plants ranged from 0.0003 to 0.0163. The higher non-carcinogenic risks occurred in the eastern study area where soils and plants were underlain by human activities (Fig. 5). The lowest non-carcinogenic risk value was acquired from inhalation, followed by dermal adsorption and ingestion pathways for children and adults from soils and plants. Total non-carcinogenic risk values of soils and plants obtained in this study were less than 1.0000, illustrating low non-carcinogenic risk of REEs. The non-carcinogenic risk posed by REEs from soil and plant ingestion were relatively low in the study area by considering the low REEs concentrations in soils and plants.



**Fig. 5** Non-carcinogenic risk of REEs from soils and plants in the northeastern Qinghai-Tibet Plateau, China. (a and b), non-carcinogenic risk of REEs for adults and children from soils, respectively; (c and d), non-carcinogenic risk of REEs for adults and children from plants, respectively. *HI* is the sum of the hazard quotient of ingestion, inhalation, and dermal adsorption.

Previous studies on the health evaluation of REEs in polluted areas have shown that the non-carcinogenic risk values exceeded more than  $3.000 \times 10^{-3}$  for adults and  $1.800 \times 10^{-2}$  for children (Pagano et al, 2015; Ferreira et al., 2022), significantly higher than those values observed in plants from this study. The average non-carcinogenic risk values of La, Ce, and Nd in soils for adults was higher than  $3.000 \times 10^{-3}$ , while the non-carcinogenic risk value of Ce in soil for children was higher than  $1.800 \times 10^{-2}$ . Normally, children are more susceptible to ingesting REEs due to hand and mouth movements. In addition, children exhibited greater health risks for incomplete formation of physiologic system and low body weight (Sun et al., 2017; Li et al., 2022). Based on the results of hazard quotient in Chinese cities, total non-carcinogenic risk values higher than 0.1000 for children should be more attention (Sun et al., 2017; Guo et al., 2019).

The effect of carcinogenic risk of REEs in soils and plants of the study area on human beings were analyzed (Fig. 6; Tables S1 and S2). The total carcinogenic risk values of REEs in soils ranged from  $1.593 \times 10^{-10}$  to  $8.839 \times 10^{-10}$  (Fig. 6a), while the total carcinogenic risk values of REEs in plants ranged from  $2.976 \times 10^{-12}$  to  $1.390 \times 10^{-10}$  (Fig. 6b). Carcinogenic risk posed by REEs in the study area had similar distribution. Carcinogenic risk through ingestion, dermal

adsorption, and inhalation were all below safe level ( $<1.000\times10^{-6}$ ) in the study area, illustrating that carcinogenic risk generated through REEs to human body was acceptable in the study area.



Fig. 6 Carcinogenic risk (CR) of REEs from soils (a) and plants (b) in the northeastern Qinghai-Tibet Plateau, China

# 4 Conclusions

The distribution of REEs in soils was noticeably homogeneous although that in plants was not homogeneous.  $\Sigma$ LREEs in soils and plants account for 76% and 77% of total REEs, respectively. Most of REEs might have similar sources in soils or plants. The distribution patterns of REEs in soils and plants from the study area were generally similar.  $\Sigma$ LREEs had significant fractionation while  $\Sigma$ HREEs had moderate fractionation in soils and plants.  $\Sigma$ LREEs enriched in soils and plants. Significant negative anomaly of Eu occurred in soils and plants. Ce was not significantly anomaly in soils and had slightly positive anomaly in plants. The average BCF values followed the order of Eu>Lu>Y>Ce>Yb>Ho>Tm>Dy>La>Tb>Er>Gd>Sm>Nd>Pr>Sc. BCF values for REEs indicated that they had relatively low bioaccumulation. Eu had relatively high mobility of transplants of the study area. These characteristic parameters of REEs in soils and plants of the study area were similar to the background values in Qinghai Province of China and the background values in China. Both non-carcinogenic and carcinogenic risks posed by REEs in soils and plants were acceptable. Adults were less sensitive to REEs ingestion compared to children.

## Acknowledgements

This work was supported by the One Hundred Talents Program of Chinese Academy of Sciences (Y610061033), the Kunlun Talent Action Plan of Qinghai Province, China (E140WX42), the Science and Technology Plan Protect of Qinghai Province, China (2021-ZJ-T07), the Taishan Scholar Program of Shandong Province, China (tsqn201812116), and the Two-Hundred Talents Plan of Yantai City of Shandong Province, China. The authors would like to thank the research group of the State Key Laboratory of Marine Geology for rare earth elements analysis. The authors also thank the anonymous reviewers and editors for their suggestions on improving the manuscript.

#### References

Aide M T, Aide C. 2012. Rare earth elements: their importance in understanding soil genesis. International Scholarly Research Notices, 2012: 783876, doi: 10.5402/2012/783876.

Allajbeu S, Yushin N S, Qarri F, et al. 2016. Atmospheric deposition of rare earth elements in Albania studied by the moss biomonitoring technique, neutron activation analysis and GIS technology. Environmental Science and Pollution Research, 23: 14087–14101.

Barta C A, Sachs-Barrable K, Jia J, et al. 2007. Lanthanide containing compounds for therapeutic care in bone resorption disorders. Dalton Transactions, 21(43): 5019–5030.

Brioschi L, Steinmann M, Lucot E, et al. 2013. Transfer of rare earth elements (REE) from natural soil to plant systems:

implications for the environmental availability of anthropogenic REE. Plant and Soil, 366: 143–163.

- Chen J Q, Wang J P, Zhang Y L, et al. 2022. A study on the distribution characteristics of arsenic and mercury in the salt lake of Ayacuo, Tibet. Journal of Salt Lake Research, 30(2): 61–69. (in Chinese)
- Chen J Y, Yang R D. 2010. Analysis on REE geochemical characteristics of three types of REE-rich soil in Guizhou Province, China. Journal of Rare Earths, 28: 517–522.
- CNEMC (China National Environmental Monitoring Centre). 1989. The Background Values of Chinese Soils. Beijing: Environmental Science Press of China, 87–90. (in Chinese)
- Costa L, Johannesson K, Mirlean N, et al. 2021. Rare earth element distributions in salt marsh sediment cores reveal evidence of environmental lability during bioturbation and diagenetic processes. Chemical Geology, 584: 120503, doi: 10.1016/j.chemgeo.2021.120503.
- Cui L, Li J, Gao X Y, et al. 2022. Human health ambient water quality criteria for 13 heavy metals and health risk assessment in Taihu Lake. Frontiers of Environmental Science & Engineering, 16: 41, doi: 10.1007/s11783-021-1475-6.
- Duan X L. 2012. Research Methods of Exposure Factor and its Application in Environmental Health Risk Assessment. Beijing: Science Press. (in Chinese)
- Durn G, Perković I, Stummeyer J, et al. 2021. Differences in the behaviour of trace and rare-earth elements in oxidizing and reducing soil environments: Case study of Terra Rossa soils and Cretaceous palaeosols from the Istrian peninsula, Croatia. Chemosphere, 283: 131286, doi: 10.1016/j.chemosphere.2021.131286.
- Faiz Y, Siddioue N, Tufail M, 2012. Pollution level and health risk assessment of road dust from an expressway. Journal of Environmental Science and Health, Part A, 47: 818–829.
- Ferreira M S, Fontes M P F, Bellato C R, et al. 2021. Geochemical signatures and natural background values of rare earth elements in soils of Brazilian Amazon. Environmental Pollution, 277: 116743, doi: 10.1016/j.envpol.2021.116743.
- Ferreira M S, Fontes M P F, Lima M T W D, et al. 2022. Human health risk assessment and geochemical mobility of rare earth elements in Amazon soils. Science of the Total Environment, 806(2): 151191, doi: 10.1016/j.scitotenv.2021.151191.
- Fu J J, Zhang Q, Huang B C, et al. 2021. A review on anammox process for the treatment of antibiotic-containing wastewater: Linking effects with corresponding mechanisms. Frontiers of Environmental Science & Engineering, 15(1): 17, doi: 10.1007/s11783-020-1309-y.
- Gill L W, Babechuk M G, Kamber B S, et al. 2018. Use of trace and rare earth elements to quantify autogenic and allogenic inputs within a lowland karst network. Applied Geochemistry, 90: 101–114.
- Godwyn-Paulson P, Jonathan M P, Rodríguez-Espinosa P F, et al. 2022. Rare earth element enrichments in beach sediments from Santa Rosalia mining region, Mexico: An index-based environmental approach. Marine Pollution Bulletin, 174: 113271, doi: 10.1016/j.marpolbul.2021.113271.
- Groenenberg J E, Romkens P F A M, Comans R N J, et al. 2010. Transfer functions for solid-solution partitioning of cadmium, copper, nickel, lead and zinc in soils: derivation of relationships for free metal ion activities and validation with independent data. European Journal of Soil Science, 61: 58–73.
- Guo G H, Song B, Lei M, et al. 2019. Rare earth elements (REEs) in PM10 and associated health risk from the polymetallic mining region of Nandan County, China. Human and Ecological Risk Assessment: An International Journal, 25: 672–687.
- He B Y, Wang J P, Kong F C, et al. 2022. Analysis of environmental pollution dissipation capacity of salt lake area in the Qaidam Basin. Journal of Salt Lake Research, 30(2): 52–60. (in Chinese)
- Henderson P. 1984. Chapter 1–general geochemical properties and abundances of the rare earth elements. In: Henderson P (Ed.). Developments in Geochemistry. Amsterdam: Elsevier, 1–32.
- Hu J L, Wu F Y, Wu S C, et al. 2013. Phytoavailability and phytovariety codetermine the bioaccumulation risk of heavy metal from soils, focusing on Cd-contaminated vegetable farms around the Rearl River delta, China. Ecotoxicology and Environmental Safety, 91(2): 18–24.
- Hu Z, Haneklaus S, Sparovek G, et al. 2006. Rare earth elements in soils. Communications in Soil Science and Plant Analysis, 37(9–10): 1381–1420.
- Huang H B, Lin C Q, Yu R L, et al. 2019. Spatial distribution and source appointment of rare earth elements in paddy soils of Jiulong River Basin, Southeast China. Journal of Geochemical Exploration, 200: 213–220.
- Huang X, Zhang G C, Pan A, et al. 2016. Protection the environment and public health from rare earth mining. Earth's Future, 4: 532–535.
- Ingrid C M S, Laís A S, Vinicius F P, et al. 2022. Environmental settings of seagrass meadows control rare earth element distribution and transfer from soil to plant compartments. Science of the Total Environment, 843: 157095, doi: 10.1016/j.scitotenv.2022.157095.
- Jeelani N, Zhu Z J, Wang P H, et al. 2017. Assessment of trace metal contamination and accumulation in sediment and plants of the Suoxu River, China. Aquatic Botany, 140: 92–95.
- Khan A M, Bakar N K A, Bakar A F A, et al. 2017. Chemical speciation and bioavailability of rare earth elements (REEs) in the

ecosystem: a review. Environmental Science and Pollution Research, 24: 22764-22789.

- Li L M, Wu J, Lu J, et al. 2018. Distribution, pollution, bioaccumulation, and ecological risks of trace elements in soils of the northeastern Qinghai-Tibet Plateau. Ecotoxicology and Environmental Safety, 166: 345–353.
- Li L M, Wu J, Lu J, et al. 2022. Water quality evaluation and ecological-health risk assessment on trace elements in surface water of the northeastern Qinghai-Tibet Plateau. Ecotoxicology and Environmental Safety, 241: 113775, doi: 10.1016/j.ecoenv.2022.113775.
- Li X F, Chen Z B, Chen Z Q, et al. 2013. A human health risk assessment of rare earth elements in soil and vegetables from a mining area in Fujian Province, Southeast China. Chemosphere, 93(6): 1240–1246.
- Liang Z F, Ding Q, Wei D P, et al. 2013. Major controlling factors and predictions for cadmium transfer from the soil into spinach plants. Ecotoxicology and Environmental Safety, 93(4): 180–185.
- Lin X K, Chen X H, Li S C, et al. 2021. Sewage sludge ditch for recovering heavy metals can improve crop yield and soil environmental quality. Frontiers of Environmental Science & Engineering, 15(2): 22, doi: 10.1007/s11783-020-1314-1.
- Liu B L, Ai S W, Zhang W Y, et al. 2017. Assessment of the bioavailability, bioaccessibility and transfer of heavy metals in the soil-grain-human systems near a mining and smelting area in NW China. Science of the Total Environment, 609: 822–829.
- Liu C, Yuan M, Liu W S, et al. 2018. Element case studies: rare earth elements. In: Van der Ent A, Echevarria G, Baker A, et al., (Eds.). Agromining: Farming for Metals. Mineral Resource Reviews. Cham: Springer, 297–308.
- Liu J, Liu Z, Liu Z, et al. 2019. Source apportionment of soil PAHs and human health exposure risks quantification from sources: the Yulin National Energy and Chemical Industry Base, China as case study. Environmental Geochemistry and Health, 41: 617–632.
- Liu K, Lv J L, He W X, et al. 2015. Major factors influencing cadmium uptake from the soil into wheat plants. Ecotoxicology and Environmental Safety, 11: 207–213.
- Lu J, Lin Y C, Wu J, et al. 2021. Continental-scale spatial distribution, sources, and health risks of heavy metals in seafood: challenge for the water-food-energy nexus sustainability in coastal regions? Environmental Science and Pollution Research, 28: 63815–63828.
- Lu J, Wu J, Wang J H, 2022a. Metagenomic analysis on resistance genes in water and microplastics from a mariculture system. Frontiers of Environmental Science & Engineering, 16(1): 4, doi: 10.1007/s11783-021-1438-y.
- Lu J, Zhang Y X, Wu J, et al. 2022b. Intervention of antimicrobial peptide usage on antimicrobial resistance in aquaculture. Journal of Hazardous Materials, 427: 128154, doi: 10.1016/j.jhazmat.2021.128154.
- Malhotra N, Hsu H-S, Liang S-T, et al. 2020. An updated review of toxicity effect of the rare earth elements (REEs) on aquatic organisms. Animals, 10: 1663, doi: 10.3390/ani10091663.
- Mclennan S M. 1989. Rare earth elements in sedimentary rocks: influence of provenance and sedimentary processes. In: Lipin Bruce R, McKay G A (Eds.). Geochemistry and Mineralogy of Rare Earth Elements. Berlin: De Gruyter, 169–200.
- MEPC (Ministry of Environmental Protection of the People's Pepubic of China), 1990. Background Values of Soil Elements in China. Beijing: China Environment Science Press, 329–493.
- Meryem B, Ji H B, Gao Y, et al. 2016. Distribution of rare earth elements in agricultural soil and human body (scalp hair and urine) near smelting and mining areas of Hezhang, China. Journal of Rare Earths, 34: 1156–1167.
- Migaszewski Z M, Gałuszka A, 2015. The characteristics, occurrence, and geochemical behavior of rare earth elements in the environment: a review. Critical Reviews in Environmental Science and Technology, 45: 429–471.
- Mihajlovic J, Stark H J, Wennrich R, et al. 2015. Rare earth elements in two Luvisols developed from loess under arable and forest land use in Bavaria, Germany: Concentrations, stocks, and potential mobilities. Soil Science, 180(3): 107–123.
- Nicolas L, Charlotte C, Rémi M, et al. 2022. Implications of speciation on rare earth element toxicity: A focus on organic matter influence in Daphnia magna standard test. Environmental Pollution, 307: 119554, doi: 10.1016/j.envpol.2022.119554.
- Ou X L, Chen Z B, Chen X L, et al. 2022. Redistribution and chemical speciation of rare earth elements in an ion-adsorption rare earth tailing, Southern China. Science of the Total Environment, 821: 153369, doi: 10.1016/j.scitotenv.2022.153369.
- Pagano G, Guida M, Tommasi F, et al. 2015. Health effects and toxicity mechanisms of rare earth elements-Knowledge gaps and research prospects. Ecotoxicology and Environmental Safety, 115: 40–48.
- Pagano G. 2016. Rare Earth Elements in Human and Environmental Health (1st ed.). Singapore: Jenny Stanford.
- Pirarath R, Shivashanmugam P, Syed A, et al. 2021. Mercury removal from aqueous solution using petal-like MoS<sub>2</sub> nanosheets. Frontiers of Environmental Science & Engineering, 15(1): 15, doi: 10.1007/s11783-020-1307-0.
- Qiao W C, Li R, Tang T H, et al. 2021. Removal, distribution and plant uptake of perfluorooctane sulfonate (PFOS) in a simulated constructed wetland system. Frontiers of Environmental Science & Engineering, 15(2): 20, doi: 10.1007/s11783-020-1312-3.
- Rodríguez-Barranco M, Lacasaña M, Gil F, et al. 2014. Cadmium exposure and neuropsychological development in school children in south-western Spain. Environmental Research, 134: 66–73.
- Shen M, Chen L Y, Han W L, et al. 2018. Methods for the determination of heavy metals in indocalamus leaves after different

preservation treatment using inductively-coupled plasma mass spectrometry. Microchemical Journal, 139: 295-300.

- State Environmental Protection Administration, 2004. The Technical Specification for Soil Environmental Monitoring (HJ/T 166–2004). China Environmental Press Co., Ltd, Beijing.
- Sun G Y, Li Z G, Liu T, et al. 2017. Rare earth elements in street dust and associated health risk in a municipal industrial base of central China. Environmental Geochemistry and Health, 39: 1469–1486.
- Tang S T, Zheng C L, Chen M J, et al. 2020. Geobiochemistry characteristics of rare earth elements in soil and ground water: a case study in Baotou, China. Science Reports, 10(1): 11740, doi: 10.1038/s41598-020-68661-4.
- Tao Y, Shen L, Feng C, et al. 2022. Distribution of rare earth elements (REEs) and their roles in plant growth: a review. Environmental Pollution, 298: 118540, doi: 10.1016/j.envpol.2021.118540.

Taylor S R, McLennan S M. 1985. The Continental Crust: Its Composition and Evolution. Oxford: Blackwell, 312.

Tyler G. 2004. Rare earth elements in soil and plant systems-a review. Plant and Soil, 267(1-2): 191-206.

USEPA (United States Environmental Protection Agency). 1996. Soil screening guidance: Technical background document. In: Office of Solid Waste and Emergency Response. Washington D.C., USA.

- USEPA (United States Environmental Protection Agency). 2002. Supplemental guidance for developing soil screening, In: Office of Solid Waste and Emergency Response. Washington D.C., USA.
- USEPA (United States Environmental Protection Agency). 2004. Supplemental guidance for dermal risk assessment. In: Risk Assessment Guidance for Superfund (RAGS), Washington D.C., USA.
- Wang C F, Chen G F, Zhu Y C, et al. 2017. Assessment of leaching behavior and human bioaccessibility of rare earth elements in typical hospital waste incineration ash in China. Frontiers of Environmental Science & Engineering, 11: 5, doi: 10.1007/s11783-017-0946-2.
- Wang J, Liu S Y, Wei X D, et al. 2022. Uptake, organ distribution and health risk assessment of potentially toxic elements in crops in abandoned indigenous smelting region. Chemosphere, 292: 133321, doi: 10.1016/j.chemosphere.2021.133321.
- Wang L J, Zhang S, Gao X J. 1998. Geochemical characteristics of rare earth elements in different types of soils in China. Journal of Rare Earths, 19(1): 51–67.
- Wang L Q, Liang T. 2016. Anomalous abundance and redistribution patterns of rare earth elements in soils of a mining area in Inner Mongolia China. Environmental Science and Pollution Research, 23: 11330–11338.
- Wang L Q, Han X X, Liang T, et al. 2019. Discrimination of rare earth element geochemistry and co-occurrence in sediment from Poyang Lake, the largest freshwater lake in China. Chemosphere, 217: 851–857.
- Wu H H, Xu C B, Wang J H, et al. 2021. Health risk assessment based on source identification of heavy metals: A case study of Beiyun River, China. Ecotoxicology and Environmental Safety, 213: 112046, doi: 10.1016/j.ecoenv.2021.112046.
- Wu J, Lu J, Li L M, et al. 2018. Pollution, ecological-health risks, and sources of heavy metals in soil of the northeastern Qinghai-Tibet Plateau. Chemosphere, 201: 234–242.
- Wu J, Lu J, Li L M, et al. 2019. Distribution, pollution, and ecological risks of rare earth elements in soil of the northeastern Qinghai-Tibet Plateau. Human and Ecological Risk Assessment: An International Journal, 25(7): 1816–1831.
- Xu X D, Cao Z M, Zhang Z X, et al. 2016. Spatial distribution and pollution assessment of heavy metals in the surface sediments of the Bohai and Yellow Seas. Marine Pollution Bulletin, 110(1): 596–602.
- Yao P Z, Hu X, Liu X, et al. 2010. REE distribution characteristics in surface sediments from Western Xiamen Bay. Journal of the Chinese Society of Rare Earths, 28(4): 495–500. (in Chinese)
- Yu R L, Lin C Q, Yan Y, et al. 2019. Distribution and provenance implication of rare earth elements and Sr-Nd isotopes in surface sediments of Jiulong River, Southeast China. Journal of Soils and Sediments, 19: 1499–1510.
- Zhang D Y, Shen X Y, Ruan Q, et al. 2014. Effects of subchronic samarium exposure on the histopathological structure and apoptosis regulation in mouse testis. Environmental Toxicology and Pharmacology, 37: 505–512.

Zhang H, Feng J, Zhu W F, et al. 2000. Chromic toxicity of rare-earth elements on human beings: implications of blood biochemical indices in REE-high regions, South Jiangxi. Biological Trace Element Research, 73: 1–17.

- Zhao Y, Yu R L, Hu G R, et al. 2017. Characteristics and environmental significance of rare earth elements in PM<sub>2.5</sub> of Nanchang, China. Journal of Rare Earths, 35(1): 98–106.
- Zheng N, Liu J S, Wang Q C, et al. 2010a. Health risk assessment of heavy metal exposure to street dust in the zinc smelting district, Northest of China. Science of the Total Environment, 408: 726–733.
- Zheng N, Liu J S, Wang Q C, et al. 2010b. Heavy metals exposure of children from stairway and sidewalk dust in the smelting district, Northeast of China. Atmospheric Environment, 44: 3239–3245.
- Zheng N, Hou S N, Wang S J, et al. 2020. Health risk assessment of heavy metals in street dust around a zinc smelting plant in China based on bioavailability and bioaccessibility. Ecotoxicology and Environmental Safety, 197: 110617, doi: 10.1016/j.ecoenv.2020.110617.



# Appendix

Fig. S1 Pearson correlations among rare earth elements in soils (a) and plants (b) of the northeastern Qinghai-Tibet Plateau, China. La, lanthanum; Ce, cerium; Pr, praseodymium; Nd, neodymium; Sm, samarium; Eu, europium; Gd, gadolinium; Tb, terbium; Dy, dysprosium; Ho, holmium; Er, erbium; Tm, thulium; Yb, ytterbium; Lu, lutetium; Y, yttrium; Sc, scandium.

Table S1Non-carcinogenic and carcinogenic risk for rare earth elements (REEs) in soils of the northeasternQinghai-Tibet Plateau, China

	Non-ca	rcinogenic	risk for cl	nildren	Non-c	arcinogen	ic risk for	adults	Carcinogenic risks			
REE	Ingestion	Inhalation	Dermal adsorption	Total	Ingestion	Inhalation	Dermal adsorption	Total	Ingestion	Inhalation	Dermal adsorption	Total
La	1.02×10 <sup>-2</sup>	7.50×10 <sup>-6</sup>	2.86×10 <sup>-4</sup>	1.05×10 <sup>-2</sup>	4.37×10 <sup>-3</sup>	3.22×10 <sup>-6</sup>	1.75×10 <sup>-4</sup>	4.55×10 <sup>-3</sup>	8.00×10 <sup>-11</sup>	6.93×10 <sup>-12</sup>	2.52×10 <sup>-12</sup>	8.94×10 <sup>-11</sup>
Ce	2.09×10 <sup>-2</sup>	1.54×10 <sup>-5</sup>	5.86×10 <sup>-4</sup>	2.15×10 <sup>-2</sup>	8.97×10 <sup>-3</sup>	6.60×10 <sup>-6</sup>	3.58×10 <sup>-4</sup>	9.33×10 <sup>-3</sup>	1.64×10 <sup>-10</sup>	1.42×10 <sup>-11</sup>	5.18×10 <sup>-12</sup>	1.83×10 <sup>-10</sup>
Pr	2.41×10 <sup>-3</sup>	1.77×10 <sup>-6</sup>	6.74×10 <sup>-5</sup>	2.48×10 <sup>-3</sup>	1.03×10 <sup>-3</sup>	7.58×10 <sup>-7</sup>	4.11×10 <sup>-5</sup>	1.07×10 <sup>-3</sup>	1.89×10 <sup>-11</sup>	1.63×10 <sup>-12</sup>	5.95×10 <sup>-13</sup>	2.11×10 <sup>-11</sup>
Nd	8.86×10 <sup>-3</sup>	6.52×10 <sup>-6</sup>	2.48×10 <sup>-4</sup>	9.12×10 <sup>-3</sup>	3.80×10 <sup>-3</sup>	2.79×10 <sup>-6</sup>	1.52×10 <sup>-4</sup>	3.95×10 <sup>-3</sup>	6.94×10 <sup>-11</sup>	6.01×10 <sup>-12</sup>	2.19×10 <sup>-12</sup>	7.76×10 <sup>-11</sup>
Sm	1.70×10 <sup>-3</sup>	1.25×10 <sup>-6</sup>	4.77×10 <sup>-5</sup>	1.75×10 <sup>-3</sup>	7.30×10 <sup>-4</sup>	5.36×10 <sup>-7</sup>	2.91×10 <sup>-5</sup>	7.59×10 <sup>-4</sup>	1.33×10 <sup>-11</sup>	1.16×10 <sup>-12</sup>	4.21×10 <sup>-13</sup>	1.49×10 <sup>-11</sup>
Eu	3.64×10 <sup>-4</sup>	2.68×10 <sup>-7</sup>	1.02×10 <sup>-5</sup>	3.75×10 <sup>-4</sup>	1.56×10 <sup>-4</sup>	1.15×10 <sup>-7</sup>	6.23×10 <sup>-6</sup>	1.62×10 <sup>-4</sup>	2.85×10 <sup>-12</sup>	2.47×10 <sup>-13</sup>	9.01×10 <sup>-14</sup>	3.19×10 <sup>-12</sup>
Gd	1.68×10 <sup>-3</sup>	1.23×10 <sup>-6</sup>	4.70×10 <sup>-5</sup>	1.73×10 <sup>-3</sup>	7.19×10 <sup>-4</sup>	5.29×10 <sup>-7</sup>	2.87×10 <sup>-5</sup>	7.48×10 <sup>-4</sup>	1.31×10 <sup>-11</sup>	1.14×10 <sup>-12</sup>	4.15×10 <sup>-13</sup>	1.47×10 <sup>-11</sup>
Tb	2.23×10 <sup>-4</sup>	1.64×10 <sup>-7</sup>	6.23×10 <sup>-6</sup>	2.29×10 <sup>-4</sup>	9.54×10 <sup>-5</sup>	7.01×10 <sup>-8</sup>	3.81×10 <sup>-6</sup>	9.93×10 <sup>-5</sup>	1.74×10 <sup>-12</sup>	1.51×10 <sup>-13</sup>	5.51×10 <sup>-14</sup>	1.95×10 <sup>-12</sup>
Dy	1.17×10 <sup>-3</sup>	8.60×10 <sup>-7</sup>	3.27×10 <sup>-5</sup>	1.20×10 <sup>-3</sup>	5.01×10 <sup>-4</sup>	3.69×10 <sup>-7</sup>	2.00×10 <sup>-5</sup>	5.22×10 <sup>-4</sup>	9.17×10 <sup>-12</sup>	7.94×10 <sup>-13</sup>	2.89×10 <sup>-13</sup>	1.02×10 <sup>-11</sup>
Но	2.26×10 <sup>-4</sup>	1.66×10 <sup>-7</sup>	6.34×10 <sup>-6</sup>	2.33×10 <sup>-4</sup>	9.70×10 <sup>-5</sup>	7.13×10 <sup>-8</sup>	3.87×10 <sup>-6</sup>	1.01×10 <sup>-4</sup>	1.77×10 <sup>-12</sup>	1.54×10 <sup>-13</sup>	5.60×10 <sup>-14</sup>	1.98×10 <sup>-12</sup>
Er	7.19×10 <sup>-4</sup>	5.29×10 <sup>-7</sup>	2.01×10 <sup>-5</sup>	7.40×10 <sup>-4</sup>	3.08×10 <sup>-4</sup>	2.27×10 <sup>-7</sup>	1.23×10 <sup>-5</sup>	3.21×10 <sup>-4</sup>	5.64×10 <sup>-12</sup>	4.88×10 <sup>-13</sup>	1.78×10 <sup>-13</sup>	6.30×10 <sup>-12</sup>
Tm	9.08×10 <sup>-5</sup>	6.68×10 <sup>-8</sup>	2.54×10 <sup>-6</sup>	9.34×10 <sup>-5</sup>	3.89×10 <sup>-5</sup>	2.86×10 <sup>-8</sup>	1.55×10 <sup>-6</sup>	4.05×10 <sup>-5</sup>	7.12×10 <sup>-13</sup>	6.16×10 <sup>-14</sup>	2.25×10 <sup>-14</sup>	7.96×10 <sup>-13</sup>
Yb	5.99×10 <sup>-4</sup>	4.40×10 <sup>-7</sup>	1.68×10 <sup>-5</sup>	6.16×10 <sup>-4</sup>	2.57×10 <sup>-4</sup>	1.89×10 <sup>-7</sup>	1.02×10 <sup>-5</sup>	2.67×10 <sup>-4</sup>	4.69×10 <sup>-12</sup>	4.06×10 <sup>-13</sup>	1.48×10 <sup>-13</sup>	5.25×10 <sup>-12</sup>
Lu	8.87×10 <sup>-5</sup>	6.52×10 <sup>-8</sup>	2.48×10 <sup>-6</sup>	9.12×10 <sup>-5</sup>	3.80×10 <sup>-5</sup>	2.79×10 <sup>-8</sup>	1.52×10 <sup>-6</sup>	3.95×10 <sup>-5</sup>	6.95×10 <sup>-13</sup>	6.02×10 <sup>-14</sup>	2.19×10 <sup>-14</sup>	7.77×10 <sup>-13</sup>
Y	5.99×10 <sup>-3</sup>	4.40×10 <sup>-6</sup>	1.68×10 <sup>-4</sup>	6.16×10 <sup>-3</sup>	2.57×10 <sup>-3</sup>	1.89×10 <sup>-6</sup>	1.02×10 <sup>-4</sup>	2.67×10 <sup>-3</sup>	4.69×10 <sup>-11</sup>	4.06×10 <sup>-12</sup>	1.48×10 <sup>-12</sup>	5.24×10 <sup>-11</sup>
Sc	3.45×10 <sup>-3</sup>	2.54×10 <sup>-6</sup>	9.67×10 <sup>-5</sup>	3.55×10 <sup>-3</sup>	1.48×10 <sup>-3</sup>	1.09×10 <sup>-6</sup>	5.91×10 <sup>-5</sup>	1.54×10 <sup>-3</sup>	2.71×10 <sup>-11</sup>	2.34×10 <sup>-12</sup>	8.55×10 <sup>-13</sup>	3.03×10 <sup>-11</sup>
Total	5.87×10 <sup>-2</sup>	4.32×10 <sup>-5</sup>	1.64×10 <sup>-3</sup>	6.04×10 <sup>-2</sup>	2.52×10 <sup>-2</sup>	1.85×10 <sup>-5</sup>	1.00×10 <sup>-3</sup>	2.62×10 <sup>-2</sup>	4.60×10 <sup>-10</sup>	3.98×10 <sup>-11</sup>	1.45×10 <sup>-11</sup>	5.14×10 <sup>-10</sup>

Note: REE, rare earth element; La, lanthanum; Ce, cerium; Pr, praseodymium; Nd, neodymium; Sm, samarium; Eu, europium; Gd, gadolinium; Tb, terbium; Dy, dysprosium; Ho, holmium; Er, erbium; Tm, thulium; Yb, ytterbium; Lu, lutetium; Y, yttrium; Sc, scandium.

	Non-ca	rcinogenio	risk for c	hildren	Non-c	arcinogen	ic risk for	adults	Carcinogenic risks				
REE	Ingestion	Inhalation	Dermal adsorption	Total	Ingestion	Inhalation	Dermal adsorption	Total	Ingestion	Inhalation	Dermal adsorption	Total	
La	5.80×10 <sup>-4</sup>	4.27×10 <sup>-7</sup>	1.62×10 <sup>-5</sup>	5.97×10 <sup>-4</sup>	2.49×10 <sup>-4</sup>	1.83×10 <sup>-7</sup>	9.92×10 <sup>-6</sup>	2.59×10 <sup>-4</sup>	4.55×10 <sup>-12</sup>	3.94×10 <sup>-13</sup>	1.44×10 <sup>-13</sup>	5.08×10 <sup>-12</sup>	
Ce	1.24×10 <sup>-3</sup>	9.14×10 <sup>-7</sup>	3.48×10 <sup>-5</sup>	1.28×10 <sup>-3</sup>	5.33×10 <sup>-4</sup>	3.92×10 <sup>-7</sup>	2.13×10 <sup>-5</sup>	5.54×10 <sup>-4</sup>	9.74×10 <sup>-12</sup>	8.44×10 <sup>-13</sup>	3.08×10 <sup>-13</sup>	1.09×10 <sup>-11</sup>	
Pr	1.28×10 <sup>-4</sup>	9.38×10 <sup>-8</sup>	3.57×10 <sup>-6</sup>	1.31×10 <sup>-4</sup>	5.47×10 <sup>-5</sup>	4.02×10 <sup>-8</sup>	3.57×10 <sup>-6</sup>	5.83×10 <sup>-5</sup>	1.00×10 <sup>-12</sup>	8.66×10 <sup>-14</sup>	3.16×10 <sup>-14</sup>	1.12×10 <sup>-12</sup>	
Nd	4.71×10 <sup>-4</sup>	3.46×10 <sup>-7</sup>	1.32×10 <sup>-5</sup>	4.84×10 <sup>-4</sup>	2.02×10 <sup>-4</sup>	1.48×10 <sup>-7</sup>	8.05×10 <sup>-6</sup>	2.10×10 <sup>-4</sup>	3.69×10 <sup>-12</sup>	3.19×10 <sup>-13</sup>	1.16×10 <sup>-13</sup>	4.12×10 <sup>-12</sup>	
Sm	9.25×10 <sup>-5</sup>	6.80×10 <sup>-8</sup>	2.59×10 <sup>-6</sup>	9.52×10 <sup>-5</sup>	3.96×10 <sup>-5</sup>	2.91×10 <sup>-8</sup>	1.58×10 <sup>-6</sup>	4.13×10 <sup>-5</sup>	7.25×10 <sup>-13</sup>	6.28×10 <sup>-14</sup>	2.29×10 <sup>-14</sup>	8.11×10 <sup>-13</sup>	
Eu	2.22×10 <sup>-5</sup>	1.63×10 <sup>-8</sup>	6.21×10 <sup>-7</sup>	2.28×10 <sup>-5</sup>	9.50×10 <sup>-6</sup>	6.98×10 <sup>-9</sup>	3.79×10 <sup>-7</sup>	9.88×10 <sup>-6</sup>	1.74×10 <sup>-13</sup>	1.50×10 <sup>-14</sup>	5.48×10 <sup>-15</sup>	1.94×10 <sup>-13</sup>	
Gd	9.24×10 <sup>-5</sup>	6.80×10 <sup>-8</sup>	2.59×10 <sup>-6</sup>	9.51×10 <sup>-5</sup>	3.96×10 <sup>-5</sup>	2.91×10 <sup>-8</sup>	1.58×10 <sup>-6</sup>	4.12×10 <sup>-5</sup>	7.24×10 <sup>-13</sup>	6.27×10 <sup>-14</sup>	2.29×10 <sup>-14</sup>	8.10×10 <sup>-13</sup>	
Tb	1.23×10 <sup>-5</sup>	9.06×10 <sup>-9</sup>	3.45×10 <sup>-7</sup>	1.27×10 <sup>-5</sup>	5.28×10 <sup>-6</sup>	3.88×10 <sup>-9</sup>	2.11×10 <sup>-7</sup>	5.49×10 <sup>-6</sup>	9.65×10 <sup>-14</sup>	8.36×10 <sup>-15</sup>	3.05×10 <sup>-15</sup>	1.08×10 <sup>-13</sup>	
Dy	6.64×10 <sup>-5</sup>	4.88×10 <sup>-8</sup>	1.86×10 <sup>-6</sup>	6.83×10 <sup>-5</sup>	2.84×10 <sup>-5</sup>	2.09×10 <sup>-8</sup>	1.13×10 <sup>-6</sup>	2.96×10 <sup>-5</sup>	5.20×10 <sup>-13</sup>	4.50×10 <sup>-14</sup>	1.64×10 <sup>-14</sup>	5.81×10 <sup>-13</sup>	
Ho	1.31×10 <sup>-5</sup>	9.65×10 <sup>-9</sup>	3.67×10 <sup>-7</sup>	1.35×10 <sup>-5</sup>	5.62×10 <sup>-6</sup>	4.13×10 <sup>-9</sup>	2.24×10 <sup>-7</sup>	5.85×10 <sup>-6</sup>	1.03×10 <sup>-13</sup>	8.90×10 <sup>-15</sup>	3.25×10 <sup>-15</sup>	1.15×10 <sup>-13</sup>	
Er	3.75×10 <sup>-5</sup>	2.76×10 <sup>-8</sup>	1.05×10 <sup>-6</sup>	3.86×10 <sup>-5</sup>	1.61×10 <sup>-5</sup>	1.18×10 <sup>-8</sup>	6.42×10 <sup>-7</sup>	1.67×10 <sup>-5</sup>	2.94×10 <sup>-13</sup>	2.55×10 <sup>-14</sup>	9.28×10 <sup>-15</sup>	3.29×10 <sup>-13</sup>	
Tm	5.26×10 <sup>-6</sup>	3.86×10 <sup>-9</sup>	1.47×10 <sup>-7</sup>	5.41×10 <sup>-6</sup>	2.25×10 <sup>-6</sup>	1.66×10 <sup>-9</sup>	8.99×10 <sup>-8</sup>	2.34×10 <sup>-6</sup>	4.12×10 <sup>-14</sup>	3.57×10 <sup>-15</sup>	1.30×10 <sup>-15</sup>	4.61×10 <sup>-14</sup>	
Yb	3.48×10 <sup>-5</sup>	2.56×10 <sup>-8</sup>	9.74×10 <sup>-7</sup>	3.58×10 <sup>-5</sup>	1.49×10 <sup>-5</sup>	1.10×10 <sup>-8</sup>	5.95×10 <sup>-7</sup>	1.55×10 <sup>-5</sup>	2.73×10 <sup>-13</sup>	2.36×10 <sup>-14</sup>	8.61×10 <sup>-15</sup>	3.05×10 <sup>-13</sup>	
Lu	5.27×10 <sup>-6</sup>	3.88×10 <sup>-9</sup>	1.48×10 <sup>-7</sup>	5.42×10 <sup>-6</sup>	2.26×10 <sup>-6</sup>	1.66×10 <sup>-9</sup>	9.01×10 <sup>-8</sup>	2.35×10 <sup>-6</sup>	4.13×10 <sup>-14</sup>	3.58×10 <sup>-15</sup>	1.30×10 <sup>-15</sup>	4.62×10 <sup>-14</sup>	
Y	3.51×10 <sup>-4</sup>	2.58×10 <sup>-7</sup>	9.82×10 <sup>-6</sup>	3.61×10 <sup>-4</sup>	1.50×10 <sup>-4</sup>	1.11×10 <sup>-7</sup>	6.00×10 <sup>-6</sup>	1.56×10 <sup>-4</sup>	2.75×10 <sup>-12</sup>	2.38×10 <sup>-13</sup>	8.68×10 <sup>-14</sup>	3.07×10 <sup>-12</sup>	
Sc	1.55×10 <sup>-4</sup>	1.14×10 <sup>-7</sup>	4.34×10 <sup>-6</sup>	1.59×10 <sup>-4</sup>	6.64×10 <sup>-5</sup>	4.88×10 <sup>-8</sup>	2.65×10 <sup>-6</sup>	6.91×10 <sup>-5</sup>	1.21×10 <sup>-12</sup>	1.05×10 <sup>-13</sup>	3.83×10 <sup>-14</sup>	1.36×10 <sup>-12</sup>	
Total	3.31×10 <sup>-3</sup>	2.43×10 <sup>-6</sup>	9.26×10 <sup>-5</sup>	3.40×10 <sup>-3</sup>	1.42×10 <sup>-3</sup>	1.04×10 <sup>-6</sup>	5.80×10 <sup>-5</sup>	1.48×10 <sup>-3</sup>	2.59×10 <sup>-11</sup>	2.25×10 <sup>-12</sup>	8.19×10 <sup>-13</sup>	2.90×10 <sup>-11</sup>	

LI Leiming et al.: Geochemical signatures and human health risk evaluation of rare...