



# Relations between the Fractional Variation of the Ionizing Continuum and C IV Broad Absorption Lines with Different Ionization Levels

Ying-Ru Lin<sup>1</sup>, Cai-Juan Pan<sup>2</sup>, and Wei-Jian Lu<sup>1</sup> 

<sup>1</sup> School of Information Engineering, Baise University, Baise 533000, China; [yingru\\_lin@qq.com](mailto:yingru_lin@qq.com), [william\\_lo@qq.com](mailto:william_lo@qq.com)

<sup>2</sup> School of Materials Science and Engineering, Baise University, Baise 533000, China

Received 2023 July 15; revised 2023 September 19; accepted 2023 October 17; published 2024 January 31

## Abstract

This paper explores the correlation between the fractional variation of the ionizing continuum and C IV broad absorption lines (BALs) with different ionization levels. Our results reveal anti-correlations between fractional variation of the continuum and fractional equivalency width (EW) variation of the C IV BALs without Al III BAL/mini-BALs at corresponding velocities, providing evidence for the widespread influence of the ionizing continuum variability on the variation of HiBALs. Conversely, for C IV BALs accompanied by Al III BAL/mini-BALs (LoBAL groups), no significant correction is detected. The absence of such a correlation does not rule out the possibility that variations in these low-ionization lines are caused by ionizing continuum variability, but rather suggests the influence of BAL saturation to some extent. This saturation effect is reflected in the distribution of the fractional EW variation, where the C IV BAL group accompanied by Al III BAL has a smaller standard deviation for the best-fitting Gaussian component than the two BAL groups without Al III BAL. However, the distribution of fractional variation of their continuum does not show any significant difference. Besides the saturation influence, another potential explanation for the lack of correlations in the LoBAL groups may be the effects of other variability mechanisms besides the ionization change, such as clouds transiting across the line of sight.

*Key words:* (galaxies:) quasars: absorption lines – (galaxies:) quasars: general – galaxies: active

## 1. Introduction

Quasar outflows have been proposed as a kind of potential active galactic nucleus (AGN) feedback mechanism, which could play a role in terminating star formation in the interstellar medium (ISM) and regulating accretion onto the supermassive black hole (SMBH; Fabian 2012; King & Pounds 2015). In rest-frame UV spectra of quasars, outflows typically appear as broad absorption lines (BALs), with absorption widths greater than  $2000 \text{ km s}^{-1}$  (Weymann et al. 1991). “Mini-BALs” are defined as absorption troughs with full widths at half minimum from a few hundred to  $2000 \text{ km s}^{-1}$  (Hamann & Sabra 2004). Most BALs have only high-ionization transitions, such as N V, Si IV, and C IV (referred to as HiBALs, as mentioned by Weymann et al. 1991). A smaller fraction, about 15% in BAL quasars, also exhibits absorption from low-ionization transitions, including Al II, Al III and Mg II (known as LoBALs, as reported by Voit et al. 1993 and Gibson et al. 2009). Another subgroup, known as FeLoBALs, represents a small subset of LoBALs that exhibit absorption of Fe II or Fe III (Wampler et al. 1995).

BAL variability can serve as a means to assess BAL structures, wind locations, and dynamics, and constrain the physical mechanisms responsible for AGN outflows. Time variability in individual sources or samples with multi-epoch observations has been reported for HiBALs (e.g., Capellupo

et al. 2011, 2012; Filiz Ak et al. 2013; Yi et al. 2019a, 2019b; He et al. 2019), for LoBALs (Zhang et al. 2011; Vivek et al. 2014; McGraw et al. 2017; Sun et al. 2017; Yi et al. 2019a; Yi & Timlin 2021), and for FeLoBALs (Vivek et al. 2012; McGraw et al. 2015; Zhang et al. 2015; Rafiee et al. 2016; Stern et al. 2017). The observed BAL variability can usually be explained by variations in the ionization parameter (e.g., Barlow 1994; Crenshaw et al. 2003; Filiz Ak et al. 2013; Wang et al. 2015; He et al. 2017; Lu & Lin 2018; Lu et al. 2018; Vivek 2019), clouds transiting across the line of sight (e.g., Lundgren et al. 2007; Gibson et al. 2008; Hamann et al. 2008; Krongold et al. 2010; Hall et al. 2011; Filiz Ak et al. 2012; Vivek et al. 2012; Capellupo et al. 2014; Shi et al. 2016; Vivek et al. 2016; Capellupo et al. 2017), or a combination of them (e.g., Capellupo et al. 2012; Vivek et al. 2014). One effective way to determine the variation mechanism of BAL in quasar sample is to investigate the correlation between variability in absorption lines and that of the continuum. Early study based on relatively small quasar samples reported no detectable correlation between changes in the BAL equivalency width (EW) and the continuum flux (Gibson et al. 2008; Vivek et al. 2014; Wildy et al. 2014). However, recent studies have found anti-correlations between the fractional variation of the continuum and that of both C IV and Si IV BAL troughs in several BAL quasar samples (Lu & Lin 2018; Lu et al. 2018;

Vivek 2019). These anti-correlations reveal the widespread influence of ionizing continuum variability on BALs, providing evidence for photoionization-driven BAL variability.

The question of whether these correlations extend to lower and higher ionization levels is still under debate. Vivek et al. (2014) found no clear correlation between absorption line and the continuum flux variabilities from a quasar sample of 22 low-ionization quasars that include Mg II and Al III BALs. However, Vivek et al. (2014) pointed out that targets with large absorption line variability also present large variability in their light curves, and they concluded that the observed variability can be well explained by the combination of two mechanisms: one is the continuum variations, and another is the clouds transiting across our line of sight.

In typical disk-wind models (e.g., Murray et al. 1995; Proga et al. 2000; Higginbottom et al. 2013), BALs with different ionization levels can be explained by viewing angle effect. More concretely, lower degrees of ionization are observed along lines of sight closer to the accretion disk plane (Hamann & Sabra 2004; Baskin et al. 2013, and Figure 16 of Filiz Ak et al. 2014). This view is supported by observations as well as simulation studies. For example, Filiz Ak et al. (2014) studied the variability characteristics of C IV BAL troughs and whether they also contained accompanying Si IV and/or Al III BAL troughs. The measurements that they conducted demonstrate the correlation between the changes in ionization level, kinematics, and column density, which is consistent with the prediction of accretion-disk wind models. For example, low ionization absorption features in line of sight typically exhibit deeper and broader C IV BAL troughs, but exhibit smaller minimum velocities and less variation. Recent numerical simulations have also shown that when low ionization absorption features presented, the BALs are wider and deeper, whereas high ionization species exhibit higher blue-edge velocities compared to the low ionization lines (Matthews et al. 2016).

Filiz Ak et al. (2014) have reported the correlated changes of BAL with different ionization levels. Building upon their work, this paper aims to explore the correlation between the fractional variation of the ionizing continuum and C IV BALs with different ionization levels. The rest of this paper is organized as follows. Section 2 presents the data preparation and measurements of BALs and spectra. Section 3 contains the results, discussions, and conclusions.

## 2. Data Preparation and Measurements

The initial sample of this paper containing 671 quasars with at least two-epoch observations was selected by Filiz Ak et al. (2014) from SDSS-I/II/III. The selection criteria for these 671 targets were described in Section 2 of Filiz Ak et al. (2014). In their study, Filiz Ak et al. (2014) identified C IV, Si IV and Al III BAL/mini-BALs in the spectra of these quasars and

**Table 1**  
Number and Spearman Rank Correlation Analysis Results of Nine Groups of BALs

BAL types	Total BAL Number <sup>a</sup>	Variation BAL Number <sup>b</sup>	$r^c$	$P^d$
C IV <sub>00</sub>	113	62	-0.46	1.97E-04
C IV <sub>s0</sub>	271	161	-0.43	1.25E-08
C IV <sub>S0</sub>	246	144	-0.26	1.33E-03
Si IV <sub>S0</sub>	246	135	-0.39	2.67E-06
C IV <sub>sa</sub>	32	16	-0.29	0.27
C IV <sub>SA</sub>	94	49	-0.26	0.07
C IV <sub>SA</sub>	95	38	-0.25	0.13
Si IV <sub>SA</sub>	95	39	-0.18	0.27
Al III <sub>SA</sub>	95	32	-0.31	0.08

**Notes.**

<sup>a</sup> BAL number selected by Filiz Ak et al. (2014).


<sup>b</sup> BAL number selected in this article.

<sup>c,d</sup> The Spearman coefficient of correlation ( $r$ ) and  $P$ -value ( $P$ ) are given in bold where  $P < 0.01$ . For the C IV<sub>00</sub>, C IV<sub>s0</sub>, C IV<sub>S0</sub>, C IV<sub>sa</sub>, C IV<sub>SA</sub>, and C IV<sub>SA</sub> groups, the  $\Delta EW / \langle EW \rangle$  is measured for C IV BAL troughs. For the Si IV<sub>S0</sub> and Si IV<sub>SA</sub> groups, the  $\Delta EW / \langle EW \rangle$  is measured for Si IV BAL troughs. For the Al III<sub>SA</sub> group, the  $\Delta EW / \langle EW \rangle$  is measured for Al III BAL troughs.

classified the C IV BALs into six groups according to whether there are Si IV and/or Al III BALs/mini-BALs at corresponding BAL regions. Specifically, “C IV<sub>00</sub>” represents C IV BALs without accompanying Si IV and Al III BALs at corresponding velocities during both epochs; “C IV<sub>s0</sub>” represents C IV BALs with a Si IV BAL at corresponding velocities in one or both epochs, but no corresponding Al III BAL; “C IV<sub>SA</sub>” represents C IV BALs with both accompanying Si IV and Al III BALs in either epoch; “C IV<sub>s0</sub>” represents C IV BALs with a Si IV mini-BAL, but no accompanying Al III BAL/mini-BAL is detected; “C IV<sub>sa</sub>” represents C IV BALs with both accompanying Si IV and Al III mini-BALs, “C IV<sub>SA</sub>” represents C IV BALs with a Si IV BAL as well as an Al III mini-BAL. In addition, Filiz Ak et al. (2014) defined the corresponding Si IV BALs of C IV<sub>S0</sub> as “Si IV<sub>S0</sub>,” the corresponding Si IV and Al III BALs of C IV<sub>SA</sub> as “Si IV<sub>SA</sub>” and “Al III<sub>SA</sub>,” respectively. It is worth noting that among these nine groups, C IV<sub>00</sub>, C IV<sub>s0</sub>, C IV<sub>S0</sub> and Si IV<sub>S0</sub> belong to HiBALs, while C IV<sub>sa</sub>, C IV<sub>SA</sub>, C IV<sub>SA</sub>, Si IV<sub>SA</sub> and Al III<sub>SA</sub> belong to loBALs.

We calculated the EW variation ( $\Delta W$ ) of all these detected BAL/mini-BALs, adopting the EW values measured by Filiz et al. (2014) and applied Equation (1) from Lu et al. (2018). We then selected BALs/mini-BALs based on the criterion of  $\Delta W > 3\sigma$ . The number of BALs included in each of the nine groups from Filiz Ak et al. (2014) and those selected for this article are listed in Table 1. We then calculated the fractional EW variation ( $\Delta EW / \langle EW \rangle$ ) of our selected C IV, Si IV, and Al III BALs using Equation (2) in Lu et al. (2018).

To obtain the flux density values, we fitted a power-law continuum for each spectrum utilizing the same procedure in Lu et al. (2018). We then used the power-law continuum flux at



**Figure 1.** Fractional variation of BALs ( $\Delta EW / \langle EW \rangle$ ) as a function of the fractional variation of the continuum ( $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$ ) for nine groups of BALs. For the  $C\text{ IV}_{00}$ ,  $C\text{ IV}_{s0}$ ,  $C\text{ IV}_{S0}$ ,  $C\text{ IV}_{sa}$ ,  $C\text{ IV}_{Sa}$ , and  $C\text{ IV}_{SA}$  groups, the  $\Delta EW / \langle EW \rangle$  is measured for C IV BAL troughs. For the  $\text{Si IV}_{s0}$  and  $\text{Si IV}_{SA}$  groups, the  $\Delta EW / \langle EW \rangle$  is measured for Si IV BAL troughs. For the  $\text{Al III}_{SA}$  group, the  $\Delta EW / \langle EW \rangle$  is measured for Al III BAL troughs.


1350 Å from the two-epoch spectra of the same quasar to calculate the fractional variation of the ionizing continuum ( $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$ ), following Equation (4) in Lu et al. (2018). Figure 1 depicts the fractional variation of BALs ( $\Delta EW / \langle EW \rangle$ ) against the fractional variation of the continuum ( $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$ ) for each group, and the corresponding results of the Spearman rank correlation analysis are listed in Table 1.

### 3. Results and Discussions

As shown in Table 1, anti-correlations between the fractional flux variations of the continuum and fractional EW variations of the BALs in  $C\text{ IV}_{00}$ ,  $C\text{ IV}_{s0}$ ,  $C\text{ IV}_{S0}$  and  $\text{Si IV}_{s0}$  groups have been confirmed by the Spearman rank correlation analysis (with  $P$ -values of Spearman coefficient less than 0.01). It is worth noting that the C IV BALs with lower ionization exhibit smaller correlation coefficients. When the C IV BALs are accompanying with Al III BAL/mini-BALs ( $C\text{ IV}_{sa}$ ,  $C\text{ IV}_{Sa}$ ,  $C\text{ IV}_{SA}$ ,  $\text{Si IV}_{SA}$  and  $\text{Al III}_{SA}$ ), no significant correlation is detected. These statistical results are consistent with the expectations of the disk-wind model and provide important insights into the variation mechanism of BALs with different ionization levels.

On one hand, we confirm the anti-correlations between the fractional variations of the continuum and BALs in four groups ( $C\text{ IV}_{00}$ ,  $C\text{ IV}_{s0}$ ,  $C\text{ IV}_{S0}$  and  $\text{Si IV}_{s0}$ ) of HiBALs. Previous studies have observed anti-correlations between the fractional variations of the continuum and absorption lines in several BAL samples (Lu & Lin 2018; Lu et al. 2018; Vivek 2019) as evidence for ionization-driven BAL variation. However, these studies did not compare the impact of different ionization states of BAL on the correlation. Our study expands on this by finding anti-correlations between the fractional flux variations of the continuum and fractional EW variations for four BAL groups without Al III BAL/mini-BAL, covering a wider ionization-potential range. This finding provides further evidence of the widespread impact of ionizing continuum variability on the variation of HiBALs (e.g., Weymann et al. 1991). Additionally, in accordance with the typical disk-wind model (e.g., Murray et al. 1995; Proga et al. 2000; Higginbottom et al. 2013), if different groups of BALs represent different viewing inclinations, the anti-correlations observed in our study provide evidence for the ubiquitous effects of ionizing continuum variability across a wide range of viewing inclinations.

On the other hand, we found no significant correlations between the fractional flux variations of the continuum and



**Figure 2.**  $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$  and  $\Delta EW / \langle EW \rangle$  distributions for the BALs in our CIV<sub>00</sub> (dotted-dashed blue), CIV<sub>S0</sub> (dashed green), and CIV<sub>SA</sub> (solid red) groups.

fractional EW variations for the five BAL groups with Al III BAL/mini-BAL (CIV<sub>sa</sub>, CIV<sub>SA</sub>, CIV<sub>Sa</sub>, Si IV<sub>SA</sub> and Al III<sub>SA</sub>), with  $P$ -values of Spearman coefficient greater than 0.01. We speculate that the lack of correlations in these groups may be due to the effects of BAL saturation or other variability mechanisms besides the ionization change, such as clouds transiting across the line of sight.

The observations of BAL profiles suffering from saturation have been reported in several previous studies (e.g., Arav 1997; Hamann 1998; Arav et al. 2001; Leighly et al. 2009; Borguet et al. 2012; Xu et al. 2018). For instance, Filiz Ak et al. (2014) have shown that when BAL troughs from lower ionization transitions are present, CIV troughs tend to be stronger and wider (see their Figure 5) and exhibit more saturation, but less fractional EW variation (see their Figure 11). This suggests that CIV troughs might experience more saturation when BAL troughs from lower ionization transitions are present. Hamann et al. (2019) also obtained empirical results showing that compared to the HiBALs, the CIV BALs in the composite spectrum of the LoBALs are deeper and wider (see their Figure 5). Using two sub-samples differing on the absorption trough depth, Vivek (2019) found that the shallow trough sub-sample exhibits an even stronger correlation while the deep trough sub-sample appears to be no correlation between the BAL variability and the continuum variability (see their Table 3). These results point out that the saturation of the BALs has a considerable effect in the correlation between the BAL variability and the continuum variability. The saturation can also be inferred from our Figure 2, which shows that the  $\Delta EW / \langle EW \rangle$  distribution of the CIV<sub>SA</sub> group has a smaller standard deviation ( $\sigma$ ) for the best-fitting Gaussian component ( $\sigma = 0.213$ ) compared to the CIV<sub>00</sub> ( $\sigma = 0.390$ ) and CIV<sub>S0</sub> ( $\sigma = 0.357$ ) groups. This is because for saturated troughs, the fractional EW variation measurements can only reflect the lower limits of the true variations in optical depth and column density, thus weaken the correlation between the absorption line variability and continuum variability. However, the  $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$  distribution of the CIV<sub>SA</sub> does not show any significant difference compared to the CIV<sub>00</sub> and CIV<sub>S0</sub> groups

(according to the K-S test results, with  $P > 0.026$ ). Considering the significant impact of line-saturation in the CIV<sub>SA</sub> group, we speculate that the effects of ionizing continuum variability may also exist in the BALs that accompany Al III BAL/mini-BALs. In other words, the ubiquitous effects of ionizing continuum variability may extend to even lower viewing inclinations.

Another potential explanation for the lack of correlations in our LoBAL groups could be the transiting of the clouds across our line of sight. Transiting scenario has been reported to be a feasible cause of the absorption-line variability (e.g., Hamann et al. 2008; Capellupo et al. 2017). For example, if a highly saturated CIV BAL exhibits clear variability, it provides evidence in favor of the transiting scenario. This is because a highly saturated CIV BAL is not easily affected by small changes in ionization, but can easily exhibit variations due to the transiting of outflows (e.g., McGraw et al. 2018). The presence of a PV BAL can serve as a detector for the highly saturated CIV BAL in the corresponding velocity (e.g., Capellupo et al. 2014, 2017; McGraw et al. 2018). To put further constraints on the origin of variability of our LoBAL subsamples, it is necessary to check individual spectra, which is beyond the scope of this paper.

In summary, we have discovered anti-correlations between the fractional variation of the continuum and fractional EW variation of the CIV BALs in four HiBAL groups, while no significant correlation was found in five LoBAL groups. The anti-correlations presented in four HiBAL groups demonstrates the widespread impact of ionizing continuum variability on the variation of outflow absorption. The lack of correlations in five LoBAL groups does not rule out the possibility that variations in these low-ionization lines are caused by ionizing continuum variability, but rather suggests the influence of BAL saturation to some extent. We have tested the  $\Delta EW / \langle EW \rangle$  and  $\Delta F_{\text{cont}} / \langle F_{\text{cont}} \rangle$  distributions of the CIV<sub>00</sub>, CIV<sub>S0</sub>, and CIV<sub>SA</sub> groups (Figure 2) to reveal the saturation influence. Another potential explanation for the lack of correlations in the LoBAL groups may be the effects of other variability mechanisms besides the ionization change, such as clouds transiting across the line of sight.

## Acknowledgments

We wish to acknowledge the reviewer for the valuable comments on this paper. We also thank Filiz Ak et al. for making their data public. This research was supported by the Guangxi Natural Science Foundation (No. 2021GXNSFBA220044), the National Natural Science Foundation of China (No. 11903002), and the Research Project of Baise University (No. 2019KN04). All observational data that support the findings of this study are available from the corresponding author (Wei-Jian Lu: [william\\_lo@qq.com](mailto:william_lo@qq.com)) on request.

Funding for the Sloan Digital Sky Survey IV was provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS website is <http://www.sdss.org/>.

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU)/University of Tokyo, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), MaxPlanck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional/MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

## ORCID iDs

Wei-Jian Lu  <https://orcid.org/0000-0002-1185-4146>

## References

Arav, N. 1997, in ASP Conf. Ser. 128, Mass Ejection from Active Galactic Nuclei, ed. N. Arav, I. Shlosman, & R. J. Weymann (San Francisco, CA: ASP), 208

Arav, N., de Kool, M., Korista, K. T., et al. 2001, *ApJ*, 561, 118

Barlow, T. A. 1994, *PASP*, 106, 548

Baskin, A., Laor, A., & Hamann, F. 2013, *MNRAS*, 432, 1525

Borguet, B. C. J., Edmonds, D., Arav, N., Benn, C., & Chamberlain, C. 2012, *ApJ*, 758, 69

Capellupo, D. M., Hamann, F., & Barlow, T. A. 2014, *MNRAS*, 444, 1893

Capellupo, D. M., Hamann, F., Herbst, H., et al. 2017, *MNRAS*, 469, 323

Capellupo, D. M., Hamann, F., Shields, J. C., Rodríguez Hidalgo, P., & Barlow, T. A. 2011, *MNRAS*, 413, 908

Capellupo, D. M., Hamann, F., Shields, J. C., Rodríguez Hidalgo, P., & Barlow, T. A. 2012, *MNRAS*, 422, 3249

Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, *ARA&A*, 41, 117

Fabian, A. C. 2012, *ARA&A*, 50, 455

Filiz Ak, N., Brandt, W. N., Hall, P. B., et al. 2012, *ApJ*, 757, 114

Filiz Ak, N., Brandt, W. N., Hall, P. B., et al. 2013, *ApJ*, 777, 168

Filiz Ak, N., Brandt, W. N., Hall, P. B., et al. 2014, *ApJ*, 791, 88

Gibson, R. R., Brandt, W. N., Schneider, D. P., & Gallagher, S. C. 2008, *ApJ*, 675, 985

Gibson, R. R., Jiang, L., Brandt, W. N., et al. 2009, *ApJ*, 692, 758

Hall, P. B., Anosov, K., White, R. L., et al. 2011, *MNRAS*, 411, 2653

Hamann, F. 1998, *ApJ*, 500, 798

Hamann, F., Herbst, H., Paris, I., & Capellupo, D. 2019, *MNRAS*, 483, 1808

Hamann, F., Kaplan, K. F., Rodríguez Hidalgo, P., Prochaska, J. X., & Herbert-Fort, S. 2008, *MNRAS*, 391, L39

Hamann, F., & Sabra, B. 2004, in ASP Conf. Ser. 311, AGN Physics with the Sloan Digital Sky Survey, ed. G. T. Richards & P. B. Hall (San Francisco, CA: ASP), 203

He, Z., Wang, T., Liu, G., et al. 2019, *NatAs*, 3, 265

He, Z., Wang, T., Zhou, H., et al. 2017, *ApJS*, 229, 22

Higginbottom, N., Knigge, C., Long, K. S., Sim, S. A., & Matthews, J. H. 2013, *MNRAS*, 436, 1390

King, A., & Pounds, K. 2015, *ARA&A*, 53, 115

Krongold, Y., Binette, L., & Hernández-Ibarra, F. 2010, *ApJL*, 724, L203

Leighly, K. M., Hamann, F., Casebeer, D. A., & Grupe, D. 2009, *ApJ*, 701, 176

Lu, W.-J., & Lin, Y.-R. 2018, *ApJ*, 862, 46

Lu, W.-J., Lin, Y.-R., & Qin, Y.-P. 2018, *MNRAS*, 473, L106

Lundgren, B. F., Wilhite, B. C., Brunner, R. J., et al. 2007, *ApJ*, 656, 73

Matthews, J. H., Knigge, C., Long, K. S., et al. 2016, *MNRAS*, 458, 293

McGraw, S. M., Brandt, W. N., Grier, C. J., et al. 2017, *MNRAS*, 469, 3163

McGraw, S. M., Shields, J. C., Hamann, F. W., et al. 2015, *MNRAS*, 453, 1379

McGraw, S. M., Shields, J. C., Hamann, F. W., Capellupo, D. M., & Herbst, H. 2018, *MNRAS*, 475, 585

Murray, N., Chiang, J., Grossman, S. A., & Voit, G. M. 1995, *ApJ*, 451, 498

Proga, D., Stone, J. M., & Kallman, T. R. 2000, *ApJ*, 543, 686

Rafiee, A., Pirkola, P., Hall, P. B., et al. 2016, *MNRAS*, 459, 2472

Shi, X.-H., Jiang, P., Wang, H.-Y., et al. 2016, *ApJ*, 829, 96

Stern, D., Graham, M. J., Arav, N., et al. 2017, *ApJ*, 839, 106

Sun, L., Zhou, H., Ji, T., et al. 2017, *ApJ*, 838, 88

Vivek, M. 2019, *MNRAS*, 486, 2379

Vivek, M., Srianand, R., & Gupta, N. 2016, *MNRAS*, 455, 136

Vivek, M., Srianand, R., Mahabal, A., & Kuriakose, V. C. 2012, *MNRAS*, 421, L107

Vivek, M., Srianand, R., Petitjean, P., et al. 2014, *MNRAS*, 440, 799

Voit, G. M., Weymann, R. J., & Korista, K. T. 1993, *ApJ*, 413, 95

Wampler, E. J., Chugai, N. N., & Petitjean, P. 1995, *ApJ*, 443, 586

Wang, T., Yang, C., Wang, H., & Ferland, G. 2015, *ApJ*, 814, 150

Weymann, R. J., Morris, S. L., Foltz, C. B., & Hewett, P. C. 1991, *ApJ*, 373, 23

Wildy, C., Goad, M. R., & Allen, J. T. 2014, *MNRAS*, 437, 1976

Xu, X., Arav, N., Miller, T., & Benn, C. 2018, *ApJ*, 858, 39

Yi, W., Brandt, W. N., Hall, P. B., et al. 2019a, *ApJS*, 242, 28

Yi, W., Vivek, M., Brandt, W. N., et al. 2019b, *ApJL*, 870, L25

Yi, W., & Timlin, J. 2021, *ApJS*, 255, 12

Zhang, S., Zhou, H., Wang, T., et al. 2015, *ApJ*, 803, 58

Zhang, S.-H., Wang, H.-Y., Zhou, H.-Y., Wang, T.-G., & Jiang, P. 2011, *RAA*, 11, 1163